

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number  
**WO 01/90108 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 487/04**

(21) International Application Number: **PCT/GB01/02158**

(22) International Filing Date: **17 May 2001 (17.05.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
0012709.2 24 May 2000 (24.05.2000) GB  
0107137.2 21 March 2001 (21.03.2001) GB

(71) Applicant (*for all designated States except US*): **MERCK SHARP & DOHME LIMITED** [GB/GB]; Hertford Road, Hoddesdon, Hertfordshire EN11 9BU (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BLACKABY, Wesley, Peter** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). **GOODACRE, Simon, Charles** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). **HALLETT, David, James** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). **JENNINGS, Andrew** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). **LEWIS, Richard, Thomas** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).

**MOORE, Kevin, William** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). **STREET, Leslie, Joseph** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).

(74) Agent: **THOMPSON, John**; Merck & Co., Inc., European Patent Dept., Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **3-PHENYL-IMIDAZO-PYRIMIDINE DERIVATIVES AS LIGANDS FOR GABA RECEPTORS**



(I)

(57) Abstract: A class of 3-phenylimidazo(1,2-*a*) pyrimidine derivatives (of Formula I, or salt or prodrug thereof: I) wherein Y represents a chemical bond, an oxygen atom, or a -NH- linkage; Z represents an optionally substituted aryl or heteroaryl group; R<sup>1</sup> represents hydrogen, hydrocarbon, a heterocyclic group, halogen, cyano, trifluoromethyl, nitro, -OR<sup>a</sup>, -SR<sup>a</sup>, -SOR<sup>a</sup>, -SO<sub>2</sub>R<sup>a</sup>, -SO<sub>2</sub>NR<sup>a</sup>R<sup>b</sup>, -NR<sup>a</sup>R<sup>b</sup>, -NR<sup>a</sup>COR<sup>b</sup>, -NR<sup>a</sup>CO<sub>2</sub>R<sup>b</sup>, -COR<sup>a</sup>, CO<sub>2</sub>R<sup>a</sup>, -CONR<sup>a</sup>R<sup>b</sup> or -CR<sup>a</sup>=NOR<sup>b</sup>; and R<sup>a</sup> and R<sup>b</sup> independently represent hydrogen, hydrocarbon or a heterocyclic group.), substituted at the *meta* position of the phenyl ring by an optionally substituted aryl or heteroaryl group which is directly attached or bridged by an oxygen atom or a -NH- linkage, are selective ligands for GABA<sub>A</sub> receptors, in particular having good affinity for the

a2 and/or a3 and/or a5 subunit thereof, and are accordingly of benefit in the treatment and/or prevention of adverse conditions of the central nervous system, including anxiety, convulsions and cognitive disorders.

WO 01/90108 A1

## 3-PHENYL-IMIDAZO-PYRIMIDINE DERIVATIVES AS LIGANDS FOR GABA RECEPTORS

5 The present invention relates to a class of substituted imidazo-  
pyrimidine derivatives and to their use in therapy. More particularly, this  
invention is concerned with imidazo[1,2-*a*]pyrimidine analogues which are  
substituted in the 3-position by a substituted phenyl ring. These  
compounds are ligands for GABA<sub>A</sub> receptors and are therefore useful in  
the therapy of deleterious mental states.

10 Receptors for the major inhibitory neurotransmitter, gamma-  
aminobutyric acid (GABA), are divided into two main classes: (1) GABA<sub>A</sub>  
receptors, which are members of the ligand-gated ion channel superfamily;  
and (2) GABA<sub>B</sub> receptors, which may be members of the G-protein linked  
receptor superfamily. Since the first cDNAs encoding individual GABA<sub>A</sub>  
15 receptor subunits were cloned the number of known members of the  
mammalian family has grown to include at least six  $\alpha$  subunits, four  $\beta$   
subunits, three  $\gamma$  subunits, one  $\delta$  subunit, one  $\epsilon$  subunit and two  $\rho$   
subunits.

20 Although knowledge of the diversity of the GABA<sub>A</sub> receptor gene  
family represents a huge step forward in our understanding of this ligand-  
gated ion channel, insight into the extent of subtype diversity is still at an  
early stage. It has been indicated that an  $\alpha$  subunit, a  $\beta$  subunit and a  $\gamma$   
subunit constitute the minimum requirement for forming a fully  
functional GABA<sub>A</sub> receptor expressed by transiently transfecting cDNAs  
25 into cells. As indicated above,  $\delta$ ,  $\epsilon$  and  $\rho$  subunits also exist, but are  
present only to a minor extent in GABA<sub>A</sub> receptor populations.

Studies of receptor size and visualisation by electron microscopy  
conclude that, like other members of the ligand-gated ion channel family,  
the native GABA<sub>A</sub> receptor exists in pentameric form. The selection of at  
30 least one  $\alpha$ , one  $\beta$  and one  $\gamma$  subunit from a repertoire of seventeen allows  
for the possible existence of more than 10,000 pentameric subunit

combinations. Moreover, this calculation overlooks the additional permutations that would be possible if the arrangement of subunits around the ion channel had no constraints (i.e. there could be 120 possible variants for a receptor composed of five different subunits).

5        Receptor subtype assemblies which do exist include, amongst many others,  $\alpha 1\beta 2\gamma 2$ ,  $\alpha 2\beta \gamma 1$ ,  $\alpha 2\beta 2/3\gamma 2$ ,  $\alpha 3\beta \gamma 2/3$ ,  $\alpha 4\beta \delta$ ,  $\alpha 5\beta 3\gamma 2/3$ ,  $\alpha 6\beta \gamma 2$  and  $\alpha 6\beta \delta$ . Subtype assemblies containing an  $\alpha 1$  subunit are present in most areas of the brain and are thought to account for over 40% of GABA<sub>A</sub> receptors in the rat. Subtype assemblies containing  $\alpha 2$  and  $\alpha 3$  subunits respectively  
10       are thought to account for about 25% and 17% of GABA<sub>A</sub> receptors in the rat. Subtype assemblies containing an  $\alpha 5$  subunit are expressed predominantly in the hippocampus and cortex and are thought to represent about 4% of GABA<sub>A</sub> receptors in the rat.

      A characteristic property of all known GABA<sub>A</sub> receptors is the  
15       presence of a number of modulatory sites, one of which is the benzodiazepine (BZ) binding site. The BZ binding site is the most explored of the GABA<sub>A</sub> receptor modulatory sites, and is the site through which anxiolytic drugs such as diazepam and temazepam exert their effect. Before the cloning of the GABA<sub>A</sub> receptor gene family, the benzodiazepine  
20       binding site was historically subdivided into two subtypes, BZ1 and BZ2, on the basis of radioligand binding studies. The BZ1 subtype has been shown to be pharmacologically equivalent to a GABA<sub>A</sub> receptor comprising the  $\alpha 1$  subunit in combination with a  $\beta$  subunit and  $\gamma 2$ . This is the most abundant GABA<sub>A</sub> receptor subtype, and is believed to represent almost  
25       half of all GABA<sub>A</sub> receptors in the brain.

      Two other major populations are the  $\alpha 2\beta \gamma 2$  and  $\alpha 3\beta \gamma 2/3$  subtypes. Together these constitute approximately a further 35% of the total GABA<sub>A</sub> receptor repertoire. Pharmacologically this combination appears to be equivalent to the BZ2 subtype as defined previously by radioligand  
30       binding, although the BZ2 subtype may also include certain  $\alpha 5$ -containing subtype assemblies. The physiological role of these subtypes has hitherto

been unclear because no sufficiently selective agonists or antagonists were known.

It is now believed that agents acting as BZ agonists at  $\alpha 1\beta\gamma 2$ ,  $\alpha 2\beta\gamma 2$  or  $\alpha 3\beta\gamma 2$  subtypes will possess desirable anxiolytic properties. Compounds which are modulators of the benzodiazepine binding site of the GABA<sub>A</sub> receptor by acting as BZ agonists are referred to hereinafter as "GABA<sub>A</sub> receptor agonists". The  $\alpha 1$ -selective GABA<sub>A</sub> receptor agonists alpidem and zolpidem are clinically prescribed as hypnotic agents, suggesting that at least some of the sedation associated with known anxiolytic drugs which act at the BZ1 binding site is mediated through GABA<sub>A</sub> receptors containing the  $\alpha 1$  subunit. Accordingly, it is considered that GABA<sub>A</sub> receptor agonists which interact more favourably with the  $\alpha 2$  and/or  $\alpha 3$  subunit than with  $\alpha 1$  will be effective in the treatment of anxiety with a reduced propensity to cause sedation. Moreover, agents which are inverse agonists of the  $\alpha 5$  subunit are likely to be beneficial in enhancing cognition, for example in subjects suffering from dementing conditions such as Alzheimer's disease. Also, agents which are antagonists or inverse agonists at  $\alpha 1$  might be employed to reverse sedation or hypnosis caused by  $\alpha 1$  agonists.

The compounds of the present invention, being selective ligands for GABA<sub>A</sub> receptors, are therefore of use in the treatment and/or prevention of a variety of disorders of the central nervous system. Such disorders include anxiety disorders, such as panic disorder with or without agoraphobia, agoraphobia without history of panic disorder, animal and other phobias including social phobias, obsessive-compulsive disorder, stress disorders including post-traumatic and acute stress disorder, and generalized or substance-induced anxiety disorder; neuroses; convulsions; migraine; depressive or bipolar disorders, for example single-episode or recurrent major depressive disorder, dysthymic disorder, bipolar I and bipolar II manic disorders, and cyclothymic disorder; psychotic disorders including schizophrenia; neurodegeneration arising from cerebral

ischemia; attention deficit hyperactivity disorder; speech disorders, including stuttering; and disorders of circadian rhythm, e.g. in subjects suffering from the effects of jet lag or shift work.

Further disorders for which selective ligands for GABA<sub>A</sub> receptors may be of benefit include pain and nociception; emesis, including acute, delayed and anticipatory emesis, in particular emesis induced by chemotherapy or radiation, as well as motion sickness, and post-operative nausea and vomiting; eating disorders including anorexia nervosa and bulimia nervosa; premenstrual syndrome; muscle spasm or spasticity, e.g. in paraplegic patients; and hearing disorders, including tinnitus and age-related hearing impairment. Selective ligands for GABA<sub>A</sub> receptors may be beneficial in enhancing cognition, for example in subjects suffering from dementing conditions such as Alzheimer's disease; and may also be effective as pre-medication prior to anaesthesia or minor procedures such as endoscopy, including gastric endoscopy.


In addition, the compounds in accordance with the present invention may be useful as radioligands in assays for detecting compounds capable of binding to the human GABA<sub>A</sub> receptor.

The present invention provides a class of imidazo-pyrimidine derivatives which possess desirable binding properties at various GABA<sub>A</sub> receptor subtypes. The compounds in accordance with the present invention have good affinity as ligands for the  $\alpha 2$  and/or  $\alpha 3$  and/or  $\alpha 5$  subunit of the human GABA<sub>A</sub> receptor. The compounds of this invention may interact more favourably with the  $\alpha 2$  and/or  $\alpha 3$  subunit than with the  $\alpha 1$  subunit; and/or may interact more favourably with the  $\alpha 5$  subunit than with the  $\alpha 1$  subunit.

The compounds of the present invention are GABA<sub>A</sub> receptor subtype ligands having a binding affinity ( $K_i$ ) for the  $\alpha 2$  and/or  $\alpha 3$  and/or  $\alpha 5$  subunit, as measured in the assay described hereinbelow, of 200 nM or less, typically of 100 nM or less, and ideally of 20 nM or less. The compounds in accordance with this invention may possess at least a 2-fold,

suitably at least a 5-fold, and advantageously at least a 10-fold, selective affinity for the  $\alpha 2$  and/or  $\alpha 3$  and/or  $\alpha 5$  subunit relative to the  $\alpha 1$  subunit. However, compounds which are not selective in terms of their binding affinity for the  $\alpha 2$  and/or  $\alpha 3$  and/or  $\alpha 5$  subunit relative to the  $\alpha 1$  subunit are also encompassed within the scope of the present invention; such compounds will desirably exhibit functional selectivity in terms of zero or weak (positive or negative) efficacy at the  $\alpha 1$  subunit and (i) a full or partial agonist profile at the  $\alpha 2$  and/or  $\alpha 3$  subunit, and/or (ii) an inverse agonist profile at the  $\alpha 5$  subunit.

The present invention provides a compound of formula I, or a salt or prodrug thereof:



(I)

wherein

Y represents a chemical bond, an oxygen atom, or a -NH- linkage;

Z represents an optionally substituted aryl or heteroaryl group;

$R^1$  represents hydrogen, hydrocarbon, a heterocyclic group, halogen, cyano, trifluoromethyl, nitro,  $-OR^a$ ,  $-SR^a$ ,  $-SOR^a$ ,  $-SO_2R^a$ ,  $-SO_2NR^aR^b$ ,

$-NR^aR^b$ ,  $-NR^aCOR^b$ ,  $-NR^aCO_2R^b$ ,  $-COR^a$ ,  $-CO_2R^a$ ,  $-CONR^aR^b$  or  $-CR^a=NOR^b$ ; and

$R^a$  and  $R^b$  independently represent hydrogen, hydrocarbon or a heterocyclic group.

The aryl or heteroaryl group Z in the compounds of formula I above may be unsubstituted, or substituted by one or more substituents.

Typically, the group Z will be unsubstituted, or substituted by one or two substituents. Suitably, the group Z is unsubstituted or monosubstituted. Illustrative substituents on the group Z include halogen, cyano, trifluoromethyl, nitro, C<sub>1-6</sub> alkoxy, amino, formyl, C<sub>2-6</sub> alkoxy carbonyl, methyloxadiazolyl, triazolyl and -CR<sup>a</sup>=NOR<sup>b</sup>, wherein R<sup>a</sup> and R<sup>b</sup> are as defined above. Typical substituents on the group Z include halogen, cyano, nitro, amino, formyl, C<sub>2-6</sub> alkoxy carbonyl and -CR<sup>a</sup>=NOR<sup>b</sup>.

For use in medicine, the salts of the compounds of formula I will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid.

Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands, e.g. quaternary ammonium salts.

The term "hydrocarbon" as used herein includes straight-chained, branched and cyclic groups containing up to 18 carbon atoms, suitably up to 15 carbon atoms, and conveniently up to 12 carbon atoms. Suitable hydrocarbon groups include C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, indanyl, aryl and aryl(C<sub>1-6</sub>)alkyl.

The expression "a heterocyclic group" as used herein includes cyclic groups containing up to 18 carbon atoms and at least one heteroatom preferably selected from oxygen, nitrogen and sulphur. The heterocyclic group suitably contains up to 15 carbon atoms and conveniently up to 12

carbon atoms, and is preferably linked through carbon. Examples of suitable heterocyclic groups include C<sub>3-7</sub> heterocycloalkyl, C<sub>3-7</sub> heterocycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl and heteroaryl(C<sub>1-6</sub>)alkyl groups.

5        Suitable alkyl groups include straight-chained and branched alkyl groups containing from 1 to 6 carbon atoms. Typical examples include methyl and ethyl groups, and straight-chained or branched propyl, butyl and pentyl groups. Particular alkyl groups are methyl, ethyl, *n*-propyl, isopropyl, isobutyl, *tert*-butyl and 2,2-dimethylpropyl. Derived expressions such as "C<sub>1-6</sub> alkoxy", "C<sub>1-6</sub> alkylamino" and "C<sub>1-6</sub> alkylsulphonyl" are to be  
10        construed accordingly.

      Suitable alkenyl groups include straight-chained and branched alkenyl groups containing from 2 to 6 carbon atoms. Typical examples include vinyl, allyl and dimethylallyl groups.

15        Suitable alkynyl groups include straight-chained and branched alkynyl groups containing from 2 to 6 carbon atoms. Typical examples include ethynyl and propargyl groups.

      Suitable cycloalkyl groups include groups containing from 3 to 7 carbon atoms. Particular cycloalkyl groups are cyclopropyl and cyclohexyl.

20        Typical examples of C<sub>3-7</sub> cycloalkyl(C<sub>1-6</sub>)alkyl groups include cyclopropylmethyl, cyclohexylmethyl and cyclohexylethyl.

      Particular indanyl groups include indan-1-yl and indan-2-yl.

      Particular aryl groups include phenyl and naphthyl, preferably phenyl.

25        Particular aryl(C<sub>1-6</sub>)alkyl groups include benzyl, phenylethyl, phenylpropyl and naphthylmethyl.

      Suitable heterocycloalkyl groups include azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl and thiomorpholinyl groups.

      A typical C<sub>3-7</sub> heterocycloalkyl(C<sub>1-6</sub>)alkyl group is morpholinylmethyl.

30        Suitable heteroaryl groups include pyridinyl, quinolinyl, isoquinolinyl, pyridazinyl, pyrimidinyl, pyrazinyl, furyl, benzofuryl,



dibenzofuryl, thienyl, benzthienyl, pyrrolyl, indolyl, pyrazolyl, indazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, benzimidazolyl, oxadiazolyl, thiadiazolyl, triazolyl and tetrazolyl groups.

The expression "heteroaryl(C<sub>1-6</sub>)alkyl" as used herein includes  
5 furylmethyl, furylethyl, thienylmethyl, thienylethyl, oxazolylmethyl, oxazolylethyl, thiazolylmethyl, thiazolylethyl, imidazolylmethyl, imidazolylethyl, oxadiazolylmethyl, oxadiazolylethyl, thiadiazolylmethyl, thiadiazolylethyl, triazolylmethyl, triazolylethyl, tetrazolylmethyl, tetrazolylethyl, pyridinylmethyl, pyridinylethyl, pyrimidinylmethyl,  
10 pyrazinylmethyl, quinolinylmethyl and isoquinolinylmethyl.

The hydrocarbon and heterocyclic groups may in turn be optionally substituted by one or more groups selected from C<sub>1-6</sub> alkyl, adamantyl, phenyl, halogen, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> aminoalkyl, trifluoromethyl, hydroxy, C<sub>1-6</sub> alkoxy, aryloxy, keto, C<sub>1-3</sub> alkylenedioxy, nitro, cyano, carboxy, C<sub>2-6</sub>  
15 alkoxycarbonyl, C<sub>2-6</sub> alkoxycarbonyl(C<sub>1-6</sub>)alkyl, C<sub>2-6</sub> alkylcarbonyloxy, arylcarbonyloxy, aminocarbonyloxy, C<sub>2-6</sub> alkylcarbonyl, arylcarbonyl, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkylsulphinyl, C<sub>1-6</sub> alkylsulphonyl, arylsulphonyl, -NR<sup>v</sup>R<sup>w</sup>, -NR<sup>v</sup>COR<sup>w</sup>, -NR<sup>v</sup>CO<sub>2</sub>R<sup>w</sup>, -NR<sup>v</sup>SO<sub>2</sub>R<sup>w</sup>, -CH<sub>2</sub>NR<sup>v</sup>SO<sub>2</sub>R<sup>w</sup>, -NHCONR<sup>v</sup>R<sup>w</sup>, -CONR<sup>v</sup>R<sup>w</sup>, -SO<sub>2</sub>NR<sup>v</sup>R<sup>w</sup> and -CH<sub>2</sub>SO<sub>2</sub>NR<sup>v</sup>R<sup>w</sup>, in which R<sup>v</sup> and R<sup>w</sup>  
20 independently represent hydrogen, C<sub>1-6</sub> alkyl, aryl or aryl(C<sub>1-6</sub>)alkyl.

The term "halogen" as used herein includes fluorine, chlorine, bromine and iodine, especially fluoro or chloro.

The present invention includes within its scope prodrugs of the compounds of formula I above. In general, such prodrugs will be  
25 functional derivatives of the compounds of formula I which are readily convertible *in vivo* into the required compound of formula I. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in *Design of Prodrugs*, ed. H. Bundgaard, Elsevier, 1985.

30 Where the compounds according to the invention have at least one asymmetric centre, they may accordingly exist as enantiomers. Where the

compounds according to the invention possess two or more asymmetric centres, they may additionally exist as diastereoisomers. It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed within the scope of the present invention.

5 In a preferred embodiment, Y represents a chemical bond.

In another embodiment, Y represents an oxygen atom.

In a further embodiment, Y represents a -NH- linkage.

Representative values for the substituent Z include phenyl, pyridinyl, thienyl, thiazolyl, imidazolyl and triazolyl, any of which groups  
10 may be optionally substituted. Typical values of Z include phenyl, pyridinyl, thienyl and thiazolyl, any of which groups may be optionally substituted. In a favoured embodiment, Z represents an optionally substituted phenyl group, in particular monosubstituted phenyl. In another embodiment, Z represents optionally substituted pyridinyl,  
15 especially pyridin-2-yl or pyridin-3-yl.

Examples of suitable substituents on the group Z include fluoro, chloro, cyano, trifluoromethyl, nitro, methoxy, amino, formyl, methoxycarbonyl, methyloxadiazolyl, triazolyl and -CH=NOH. Examples  
20 of typical substituents on the group Z include chloro, cyano, nitro, amino, formyl, methoxycarbonyl and -CH=NOH. Examples of particular substituents on the group Z include fluoro, cyano, trifluoromethyl, methoxy, methyloxadiazolyl, triazolyl and -CH=NOH; especially fluoro or cyano; and more especially cyano.

Illustrative values of Z include fluorophenyl, cyanophenyl,  
25 (cyano)(fluoro)phenyl, trifluoromethyl-phenyl, nitrophenyl, methoxyphenyl, methyloxadiazolyl-phenyl, triazolyl-phenyl, phenyl-CH=NOH, pyridinyl, (amino)(chloro)pyridinyl, cyano-pyridinyl, cyano-thienyl, formyl-thienyl, methoxycarbonyl-thienyl, thienyl-CH=NOH, thiazolyl, imidazolyl and triazolyl. Specific values of Z include  
30 cyanophenyl, nitrophenyl, pyridinyl, (amino)(chloro)pyridinyl, cyano-thienyl, formyl-thienyl, methoxycarbonyl-thienyl, thienyl-CH=NOH and

thiazolyl. Individual values of Z include fluorophenyl, cyanophenyl, (cyano)(fluoro)phenyl, trifluoromethyl-phenyl, methoxyphenyl, methyloxadiazolyl-phenyl, triazolyl-phenyl, phenyl-CH=NOH, pyridinyl, cyano-pyridinyl, thiazolyl, imidazolyl and triazolyl.

5 A particular value of Z is cyanophenyl, especially 2-cyanophenyl.

Typically, R<sup>1</sup> represents hydrogen, hydrocarbon, a heterocyclic group, halogen, cyano, trifluoromethyl, -OR<sup>a</sup>, -COR<sup>a</sup>, -CO<sub>2</sub>R<sup>a</sup> or -CR<sup>a</sup>=NOR<sup>b</sup>. Suitably, R<sup>1</sup> represents hydrocarbon, a heterocyclic group, halogen, trifluoromethyl, -OR<sup>a</sup>, -COR<sup>a</sup>, -CO<sub>2</sub>R<sup>a</sup> or -CR<sup>a</sup>=NOR<sup>b</sup>.

10 Typical values of R<sup>a</sup> include hydrogen and C<sub>1-6</sub> alkyl. Suitably, R<sup>a</sup> represents hydrogen or methyl.

Typical values of R<sup>b</sup> include hydrogen, C<sub>1-6</sub> alkyl, hydroxy(C<sub>1-6</sub>)alkyl and di(C<sub>1-6</sub>)alkylamino(C<sub>1-6</sub>)alkyl. Suitably, R<sup>b</sup> represents hydrogen, methyl, ethyl, hydroxyethyl or dimethylaminoethyl. Particular values of  
15 R<sup>b</sup> include hydrogen, hydroxyethyl and dimethylaminoethyl, especially hydrogen or dimethylaminoethyl.

Representative values of R<sup>1</sup> include hydrogen, C<sub>1-6</sub> alkyl, halo(C<sub>1-6</sub>)alkyl, dihalo(C<sub>1-6</sub>)alkyl, hydroxy(C<sub>1-6</sub>)alkyl, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, di(C<sub>1-6</sub>)alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> heterocycloalkyl(C<sub>1-6</sub>)alkyl,  
20 heteroaryl, C<sub>1-6</sub> alkyl-heteroaryl, heteroaryl(C<sub>1-6</sub>)alkyl, halogen, cyano, trifluoromethyl, C<sub>1-6</sub> alkoxy, formyl, C<sub>2-6</sub> alkylcarbonyl, C<sub>2-6</sub> alkoxycarbonyl and -CR<sup>a</sup>=NOR<sup>b</sup>, in which R<sup>a</sup> and R<sup>b</sup> are as defined above. Typical values of R<sup>1</sup> include hydrogen, C<sub>1-6</sub> alkyl, halo(C<sub>1-6</sub>)alkyl, dihalo(C<sub>1-6</sub>)alkyl, hydroxy(C<sub>1-6</sub>)alkyl, di(C<sub>1-6</sub>)alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>  
25 heterocycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl(C<sub>1-6</sub>)alkyl, cyano, trifluoromethyl, C<sub>1-6</sub> alkoxy, formyl, C<sub>2-6</sub> alkylcarbonyl, C<sub>2-6</sub> alkoxycarbonyl and -CR<sup>a</sup>=NOR<sup>b</sup>, in which R<sup>a</sup> and R<sup>b</sup> are as defined above. Illustrative values of R<sup>1</sup> include C<sub>1-6</sub> alkyl, hydroxy(C<sub>1-6</sub>)alkyl, heteroaryl, halogen, trifluoromethyl, C<sub>1-6</sub> alkoxy, formyl, C<sub>2-6</sub> alkylcarbonyl, C<sub>2-6</sub> alkoxycarbonyl  
30 and -CR<sup>a</sup>=NOR<sup>b</sup>, in which R<sup>a</sup> and R<sup>b</sup> are as defined above.

Itemised values of  $R^1$  include hydrogen, methyl, fluoromethyl, difluoromethyl, hydroxymethyl, methoxymethyl, dimethoxymethyl, hydroxyethyl (especially 1-hydroxyethyl), fluoroethyl (especially 1-fluoroethyl), difluoroethyl (especially 1,1-difluoroethyl), dimethoxyethyl (especially 1,1-dimethoxyethyl), isopropyl, hydroxypropyl (especially 2-hydroxyprop-2-yl), fluoropropyl (especially 2-fluoroprop-2-yl), *tert*-butyl, cyclopropyl, cyclobutyl, morpholinylmethyl, pyridinyl, furyl, thienyl, oxazolyl, methylthiazolyl, methyloxadiazolyl, imidazolylmethyl, triazolylmethyl, chloro, cyano, trifluoromethyl, methoxy, formyl, acetyl, methoxycarbonyl and  $-CR^2=NOR^3$ , in which  $R^2$  represents hydrogen or methyl, and  $R^3$  represents hydrogen, hydroxyethyl or dimethylaminoethyl.

Selected values of  $R^1$  include hydrogen, methyl, fluoromethyl, difluoromethyl, hydroxymethyl, dimethoxymethyl, dimethoxyethyl (especially 1,1-dimethoxyethyl), isopropyl, hydroxypropyl (especially 2-hydroxyprop-2-yl), *tert*-butyl, cyclopropyl, cyclobutyl, morpholinylmethyl, triazolylmethyl, cyano, trifluoromethyl, methoxy, formyl, acetyl, methoxycarbonyl and  $-CR^2=NOR^3$ , in which  $R^2$  and  $R^3$  are as defined above.

Individual values of  $R^1$  include hydrogen, methyl, fluoromethyl, difluoromethyl, hydroxymethyl, dimethoxymethyl, dimethoxyethyl (especially 1,1-dimethoxyethyl), isopropyl, *tert*-butyl, cyclopropyl, cyclobutyl, morpholinylmethyl, triazolylmethyl, cyano, trifluoromethyl, methoxy, formyl, acetyl, methoxycarbonyl and  $-CR^2=NOR^3$ , in which  $R^2$  and  $R^3$  are as defined above.

Specific values of  $R^1$  include methyl, hydroxymethyl, hydroxyethyl, furyl, chloro, trifluoromethyl, methoxy, formyl, acetyl, methoxycarbonyl and  $-CR^2=NOR^3$ , in which  $R^2$  and  $R^3$  are as defined above.


A particular value of  $R^1$  is  $C_{1-6}$  alkyl, especially methyl.

In one favoured embodiment,  $R^1$  represents 2-hydroxyprop-2-yl. In another favoured embodiment,  $R^1$  represents trifluoromethyl.

Suitably,  $R^2$  is hydrogen.

Suitably, R<sup>3</sup> represents hydrogen or dimethylaminoethyl, especially hydrogen.

A particular sub-class of compounds according to the invention is represented by the compounds of formula IIA, and salts and prodrugs thereof:



(IIA)

wherein

10 Z is as defined above;

R<sup>11</sup> represents hydrogen, C<sub>1-6</sub> alkyl, halo(C<sub>1-6</sub>)alkyl, dihalo(C<sub>1-6</sub>)alkyl, hydroxy(C<sub>1-6</sub>)alkyl, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, di(C<sub>1-6</sub>)alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub> heterocycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl, C<sub>1-6</sub> alkyl-heteroaryl, heteroaryl(C<sub>1-6</sub>)alkyl, halogen, cyano, trifluoromethyl, C<sub>1-6</sub> alkoxy, formyl, C<sub>2-6</sub> alkylcarbonyl, C<sub>2-6</sub> alkoxy carbonyl or -CR<sup>4</sup>=NOR<sup>5</sup>;

R<sup>4</sup> represents hydrogen or C<sub>1-6</sub> alkyl; and

R<sup>5</sup> represents hydrogen, C<sub>1-6</sub> alkyl, hydroxy(C<sub>1-6</sub>)alkyl or di(C<sub>1-6</sub>)alkylamino(C<sub>1-6</sub>)alkyl.

20 The present invention also provides a compound of formula IIA as depicted above, or a salt or prodrug thereof, wherein

R<sup>11</sup> represents C<sub>1-6</sub> alkyl, hydroxy(C<sub>1-6</sub>)alkyl, heteroaryl, halogen, trifluoromethyl, C<sub>1-6</sub> alkoxy, formyl, C<sub>2-6</sub> alkylcarbonyl, C<sub>2-6</sub> alkoxy carbonyl or -CR<sup>4</sup>=NOR<sup>5</sup>; and

Z, R<sup>4</sup> and R<sup>5</sup> are as defined above.

25 Suitably, R<sup>4</sup> represents hydrogen or methyl, especially hydrogen.

Suitably,  $R^5$  represents hydrogen, methyl, ethyl, hydroxyethyl or dimethylaminoethyl. Particular values of  $R^5$  include hydrogen, hydroxyethyl and dimethylaminoethyl. Typically,  $R^5$  represents hydrogen or dimethylaminoethyl, especially hydrogen.

5       Where  $R^{11}$  represents  $C_{3-7}$  heterocycloalkyl( $C_{1-6}$ )alkyl, this group is suitably morpholinylmethyl.

Where  $R^{11}$  represents heteroaryl, this group is suitably pyridinyl, furyl, thienyl or oxazolyl, especially furyl.

10       Where  $R^{11}$  represents  $C_{1-6}$  alkyl-heteroaryl, this group is suitably methylthiazolyl (e.g. 2-methylthiazol-5-yl) or methyloxadiazolyl (e.g. 3-methyl-[1,2,4]oxadiazol-5-yl).

Where  $R^{11}$  represents heteroaryl( $C_{1-6}$ )alkyl, this group is suitably imidazolylmethyl or triazolylmethyl.

15       Typical values of  $R^{11}$  include hydrogen,  $C_{1-6}$  alkyl, halo( $C_{1-6}$ )alkyl, dihalo( $C_{1-6}$ )alkyl, hydroxy( $C_{1-6}$ )alkyl, di( $C_{1-6}$ )alkoxy( $C_{1-6}$ )alkyl,  $C_{3-7}$  cycloalkyl,  $C_{3-7}$  heterocycloalkyl( $C_{1-6}$ )alkyl, heteroaryl( $C_{1-6}$ )alkyl, cyano, trifluoromethyl,  $C_{1-6}$  alkoxy, formyl,  $C_{2-6}$  alkylcarbonyl,  $C_{2-6}$  alkoxy carbonyl and  $-CR^4=NOR^5$ , in which  $R^4$  and  $R^5$  are as defined above.

20       Itemised values of  $R^{11}$  include hydrogen, methyl, fluoromethyl, difluoromethyl, hydroxymethyl, methoxymethyl, dimethoxymethyl, hydroxyethyl (especially 1-hydroxyethyl), fluoroethyl (especially 1-fluoroethyl), difluoroethyl (especially 1,1-difluoroethyl), dimethoxyethyl (especially 1,1-dimethoxyethyl), isopropyl, hydroxypropyl (especially 2-hydroxyprop-2-yl), fluoropropyl (especially 2-fluoroprop-2-yl), *tert*-butyl, 25       cyclopropyl, cyclobutyl, morpholinylmethyl, pyridinyl, furyl, thienyl, oxazolyl, methylthiazolyl, methyloxadiazolyl, imidazolylmethyl, triazolylmethyl, chloro, cyano, trifluoromethyl, methoxy, formyl, acetyl, methoxycarbonyl and  $-CR^2=NOR^3$ , in which  $R^2$  and  $R^3$  are as defined above.

30       Selected values of  $R^{11}$  include hydrogen, methyl, fluoromethyl, difluoromethyl, hydroxymethyl, dimethoxymethyl, dimethoxyethyl

(especially 1,1-dimethoxyethyl), isopropyl, hydroxypropyl (especially 2-hydroxyprop-2-yl), *tert*-butyl, cyclopropyl, cyclobutyl, morpholinylmethyl, triazolylmethyl, cyano, trifluoromethyl, methoxy, formyl, acetyl, methoxycarbonyl and  $-CR^2=NOR^3$ , in which  $R^2$  and  $R^3$  are as defined  
5 above.


Individual values of  $R^{11}$  include hydrogen, methyl, fluoromethyl, difluoromethyl, hydroxymethyl, dimethoxymethyl, dimethoxyethyl (especially 1,1-dimethoxyethyl), isopropyl, *tert*-butyl, cyclopropyl, cyclobutyl, morpholinylmethyl, triazolylmethyl, cyano, trifluoromethyl,  
10 methoxy, formyl, acetyl, methoxycarbonyl and  $-CR^2=NOR^3$ , in which  $R^2$  and  $R^3$  are as defined above.

Representative values of  $R^{11}$  include methyl, hydroxymethyl, hydroxyethyl, furyl, chloro, trifluoromethyl, methoxy, formyl, acetyl, methoxycarbonyl and  $-CR^2=NOR^3$ , in which  $R^2$  and  $R^3$  are as defined  
15 above.

A particular value of  $R^{11}$  is  $C_{1-6}$  alkyl, especially methyl.

In one favoured embodiment,  $R^{11}$  represents 2-hydroxyprop-2-yl. In another favoured embodiment,  $R^{11}$  represents trifluoromethyl.


One representative subset of the compounds of formula IIA above is  
20 represented by the compounds of formula IIB, and salts and prodrugs thereof:



(IIB)

25 wherein Z is as defined above.

Another representative subset of the compounds of formula IIA above is represented by the compounds of formula IIC, and salts and prodrugs thereof:



5

(IIC)

wherein

X represents CH or N;

R<sup>6</sup> represents fluoro, cyano, trifluoromethyl, methoxy,  
10 methyloxadiazolyl, triazolyl or -CR<sup>2</sup>=NOR<sup>3</sup>;

R<sup>7</sup> represents hydrogen or fluoro; and

R<sup>2</sup>, R<sup>3</sup> and R<sup>11</sup> are as defined above.

In a favoured embodiment, X represents CH. In another  
embodiment, X represents N.


15 In a particular embodiment, R<sup>6</sup> represents cyano.

In one embodiment, R<sup>7</sup> is hydrogen. In another embodiment, R<sup>7</sup> is  
fluoro.

In a specific embodiment of the compounds of formula IIC, X is CH,  
R<sup>6</sup> is cyano, and R<sup>7</sup> is hydrogen.

20 A further representative subset of the compounds of formula IIA  
above is represented by the compounds of formula IID, and salts and  
prodrugs thereof:





(IID)

wherein

W represents CH or N; and

5         $R^{11}$  is as defined above.

In one embodiment, W represents CH. In another embodiment, W represents N.

In relation to formula IID above, the substituent  $R^{11}$  favourably represents trifluoromethyl.

10        Specific compounds within the scope of the present invention include:

3'-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(imidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-trifluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

15        3'-[7-(1,1-dimethoxyethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile;

3'-(7-acetylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-isopropylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-cyclopropylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

20        3'-(7-*tert*-butylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-cyclobutylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-methoxyimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-hydroxymethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-fluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

- 3'-(7-formylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;  
3'-(7-hydroxyiminomethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;  
3-(2'-cyanobiphenyl-3-yl)imidazo[1,2-*a*]pyrimidine-7-carbonitrile;  
5 3-(2'-methoxybiphenyl-3-yl)-7-methylimidazo[1,2-*a*]pyrimidine;  
3-(2'-cyanobiphenyl-3-yl)imidazo[1,2-*a*]pyrimidine-7-carboxylic acid methyl ester;  
3'-(7-dimethoxymethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;  
10 3'-[7-([1,2,4]triazol-1-ylmethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile;  
3'-(7-difluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;  
7-methyl-3-[3-(pyridin-3-yl)phenyl]imidazo[1,2-*a*]pyrimidine;  
7-methyl-3-[3'-(5-methyl-[1,2,4]oxadiazol-3-yl)biphenyl-3-yl]imidazo[1,2-  
15 *a*]pyrimidine;  
7-methyl-3-[2'-(3-methyl-[1,2,4]oxadiazol-5-yl)biphenyl-3-yl]imidazo[1,2-*a*]pyrimidine;  
7-methyl-3-[3-(thiazol-4-yl)phenyl]imidazo[1,2-*a*]pyrimidine;  
3'-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbaldehyde oxime;  
20 3-[3-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)phenyl]pyridine-2-carbonitrile;  
7-methyl-3-[3-(pyridin-2-yl)phenyl]imidazo[1,2-*a*]pyrimidine;  
7-methyl-3-[3-(thiazol-2-yl)phenyl]imidazo[1,2-*a*]pyrimidine;  
7-methyl-3-(2'-trifluoromethylbiphenyl-3-yl)imidazo[1,2-*a*]pyrimidine;  
3-(2'-fluorobiphenyl-3-yl)-7-methylimidazo[1,2-*a*]pyrimidine;  
25 4-fluoro-3'-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;  
3-[3-(imidazol-1-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine;  
3-[3-([1,2,4]triazol-1-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine;  
3-[2'-([1,2,4]triazol-1-yl)biphenyl-3-yl]-7-trifluoromethylimidazo[1,2-  
*a*]pyrimidine;  
30 3'-[7-(morpholin-4-ylmethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile;

4-fluoro-3'-(7-trifluoromethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile;

4-fluoro-3'-[7-(2-hydroxyprop-2-yl)imidazo[1,2- $\alpha$ ]pyrimidin-3-yl]biphenyl-2-carbonitrile;

- 5 3-[3-(pyridin-3-yl)phenyl]-7-trifluoromethylimidazo[1,2- $\alpha$ ]pyrimidine;  
3-[3-([1,2,4]triazol-4-yl)phenyl]-7-trifluoromethylimidazo[1,2- $\alpha$ ]pyrimidine;  
and salts and prodrugs thereof.

Also provided by the present invention is a method for the treatment and/or prevention of anxiety which comprises administering to  
10 a patient in need of such treatment an effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof or a prodrug thereof.

Further provided by the present invention is a method for the treatment and/or prevention of convulsions (e.g. in a patient suffering from  
15 epilepsy or a related disorder) which comprises administering to a patient in need of such treatment an effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof or a prodrug thereof.

The binding affinity ( $K_i$ ) of the compounds according to the present  
20 invention for the  $\alpha 3$  subunit of the human GABA<sub>A</sub> receptor is conveniently as measured in the assay described hereinbelow. The  $\alpha 3$  subunit binding affinity ( $K_i$ ) of the anxiolytic compounds of the invention is ideally 50 nM or less, preferably 10 nM or less, and more preferably 5 nM or less.

The anxiolytic compounds according to the present invention will  
25 ideally elicit at least a 40%, preferably at least a 50%, and more preferably at least a 60%, potentiation of the GABA EC<sub>20</sub> response in stably transfected recombinant cell lines expressing the  $\alpha 3$  subunit of the human GABA<sub>A</sub> receptor. Moreover, the compounds of the invention will ideally elicit at most a 30%, preferably at most a 20%, and more preferably at  
30 most a 10%, potentiation of the GABA EC<sub>20</sub> response in stably transfected

recombinant cell lines expressing the  $\alpha 1$  subunit of the human GABA<sub>A</sub> receptor.

5 The potentiation of the GABA EC<sub>20</sub> response in stably transfected cell lines expressing the  $\alpha 3$  and  $\alpha 1$  subunits of the human GABA<sub>A</sub> receptor can conveniently be measured by procedures analogous to the protocol described in Wafford *et al.*, *Mol. Pharmacol.*, 1996, **50**, 670-678. The procedure will suitably be carried out utilising cultures of stably transfected eukaryotic cells, typically of stably transfected mouse Ltk-fibroblast cells.

10 The compounds according to the present invention may exhibit anxiolytic activity, as may be demonstrated by a positive response in the elevated plus maze and conditioned suppression of drinking tests (cf. Dawson *et al.*, *Psychopharmacology*, 1995, **121**, 109-117). Moreover, the compounds of the invention are likely to be substantially non-sedating, as  
15 may be confirmed by an appropriate result obtained from the response sensitivity (chain-pulling) test (cf. Bayley *et al.*, *J. Psychopharmacol.*, 1996, **10**, 206-213).

The compounds according to the present invention may also exhibit anticonvulsant activity. This can be demonstrated by the ability to block  
20 pentylenetetrazole-induced seizures in rats and mice, following a protocol analogous to that described by Bristow *et al.* in *J. Pharmacol. Exp. Ther.*, 1996, **279**, 492-501.

In another aspect, the present invention provides a method for the treatment and/or prevention of cognitive disorders, including dementing  
25 conditions such as Alzheimer's disease, which comprises administering to a patient in need of such treatment an effective amount of a compound of formula I as defined above or a pharmaceutically acceptable salt thereof.

Cognition enhancement can be shown by testing the compounds in the Morris watermaze as reported by McNamara and Skelton,  
30 *Psychobiology*, 1993, **21**, 101-108. Further details of relevant methodology are described in WO 96/25948.

Cognitive disorders for which the compounds of the present invention may be of benefit include delirium, dementia, amnesic disorders, and cognition deficits, including age-related memory deficits, due to traumatic injury, stroke, Parkinson's disease and Down Syndrome.

5 Any of these conditions may be attributable to substance abuse or withdrawal. Examples of dementia include dementia of the Alzheimer's type with early or late onset, and vascular dementia, any of which may be uncomplicated or accompanied by delirium, delusions or depressed mood; and dementia due to HIV disease, head trauma, Parkinson's disease or  
10 Creutzfeld-Jakob disease.

In order to elicit their behavioural effects, the compounds of the invention will ideally be brain-penetrant; in other words, these compounds will be capable of crossing the so-called "blood-brain barrier". Preferably, the compounds of the invention will be capable of exerting their beneficial  
15 therapeutic action following administration by the oral route.


The invention also provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules,  
20 sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. For preparing solid compositions such as tablets, the principal active ingredient is mixed with  
25 a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a  
30 pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active

ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described  
5 above containing from 0.1 to about 500 mg of the active ingredient of the present invention. Typical unit dosage forms contain from 1 to 100 mg, for example 1, 2, 5, 10, 25, 50 or 100 mg, of the active ingredient. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For  
10 example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A  
15 variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.


The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection  
20 include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as  
25 tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

In the treatment of neurological disorders, a suitable dosage level is about 0.01 to 250 mg/kg per day, preferably about 0.05 to 100 mg/kg per day, and especially about 0.05 to 5 mg/kg per day. The compounds may be  
30 administered on a regimen of 1 to 4 times per day.

The compounds in accordance with the present invention may be prepared by a process which comprises reacting a compound of formula III with a compound of formula IV:



(III)




(IV)

wherein Y, Z and  $R^1$  are as defined above,  $L^1$  represents a suitable leaving group, and  $M^1$  represents a boronic acid moiety  $-B(OH)_2$  or a cyclic ester thereof formed with an organic diol, e.g. pinacol, 1,3-propanediol or neopentyl glycol, or  $M^1$  represents  $-Sn(Alk)_3$  in which Alk represents a  $C_{1-6}$  alkyl group, typically *n*-butyl; in the presence of a transition metal catalyst.


The leaving group  $L^1$  is typically a halogen atom, e.g. bromo.

The transition metal catalyst of use in the reaction between compounds III and IV is suitably tetrakis(triphenylphosphine)-palladium(0). The reaction is conveniently carried out at an elevated temperature in a solvent such as *N,N*-dimethylacetamide, 1,4-dioxane or tetrahydrofuran, advantageously in the presence of potassium phosphate, copper(I) iodide or sodium carbonate. Alternatively, the transition metal catalyst employed may be dichloro[1,1'-bis(diphenylphosphino)-ferrocene]palladium(II), in which case the reaction is conveniently effected at an elevated temperature in a solvent such as *N,N*-dimethylformamide.

In an alternative procedure, the compounds according to the present invention may be prepared by a process which comprises reacting a compound of formula V with a compound of formula VI:




(V)




(VI)

wherein Y, Z, R<sup>1</sup>, L<sup>1</sup> and M<sup>1</sup> are as defined above; in the presence of a transition metal catalyst; under conditions analogous to those described above for the reaction between compounds III and IV.

5 In another procedure, the compounds according to the present invention in which Y represents a chemical bond may be prepared by a process which comprises reacting a compound of formula VII with a compound of formula VIII:



(VII)




(VIII)

wherein Z, R<sup>1</sup>, L<sup>1</sup> and M<sup>1</sup> are as defined above; in the presence of a transition metal catalyst; under conditions analogous to those described above for the reaction between compounds III and IV.


15 In the compounds of formula VII above, the leaving group L<sup>1</sup> is typically trifluoromethanesulfonyloxy (triflyloxy); or a halogen atom, e.g. bromo.

Alternatively, the compounds according to the present invention in which Y represents a chemical bond may be prepared by a process which  
20 comprises reacting a compound of formula IX with a compound of formula X:






(IX)



(X)

wherein Z, R<sup>1</sup>, L<sup>1</sup> and M<sup>1</sup> are as defined above; in the presence of a  
 5 transition metal catalyst; under conditions analogous to those described  
 above for the reaction between compounds III and IV.

In an additional procedure, the compounds according to the present  
 invention in which Y represents an oxygen atom may be prepared by a  
 process which comprises reacting a compound of formula X as defined  
 10 above with a compound of formula XI:




(XI)

wherein R<sup>1</sup> is as defined above.

15 The reaction is conveniently carried out under basic conditions, e.g.  
 using sodium hydride in a solvent such as *N,N*-dimethylformamide,  
 typically at an elevated temperature which may be in the region of 120°C.

In a further procedure, the compounds according to the present  
 invention in which Y represents a -NH- linkage may be prepared by a

process which comprises reacting a compound of formula X as defined above with a compound of formula XII:



(XII)

5


wherein R<sup>1</sup> is as defined above.

In relation to the reaction between compounds X and XII, the leaving group L<sup>1</sup> in the compounds of formula X may suitably represent fluoro.


10

The reaction between compounds X and XII is conveniently carried out by heating the reactants, typically at a temperature in the region of 120°C, in a solvent such as *N,N*-dimethylformamide.

Where M<sup>1</sup> in the intermediates of formula IV and IX above represents a cyclic ester of a boronic acid moiety -B(OH)<sub>2</sub> formed with pinacol or neopentyl glycol, the relevant compound IV or IX may be prepared by reacting bis(pinacolato)diboron or bis(neopentyl glycolato)diborane respectively with a compound of formula VIA or VIIA:



(VIA)



(VIIA)

20


wherein Y, Z and R<sup>1</sup> are as defined above, and L<sup>2</sup> represents hydroxy or a suitable leaving group; in the presence of a transition metal catalyst.

Where L<sup>2</sup> represents a leaving group, this is typically triflyloxy; or a halogen atom such as bromo.

5       The transition metal catalyst of use in the reaction between bis(pinacolato)diboron or bis(neopentyl glycolato)diborane and compound VIA or VIIA is suitably dichloro[1,1'-bis(diphenylphosphino)ferrocene]-palladium(II). The reaction is conveniently carried out at an elevated temperature in a solvent such as 1,4-dioxane, optionally in admixture with  
10       dimethylsulfoxide, typically in the presence of 1,1'-bis(diphenylphosphino)ferrocene and/or potassium acetate.

Where L<sup>1</sup>/L<sup>2</sup> in the intermediates of formula VII/VIIA above represents triflyloxy, the relevant compound VII/VIIA may be prepared by reacting the appropriate compound of formula XI as defined above with  
15       triflic anhydride, typically in the presence of pyridine. Analogous conditions may be utilised for converting an intermediate of formula VIA above wherein L<sup>2</sup> represents hydroxy into the corresponding compound of formula VI/VIA wherein L<sup>1</sup>/L<sup>2</sup> represents triflyloxy.


20       The intermediates of formula XI above may suitably be prepared from the appropriate methoxy-substituted precursor of formula XIII:



(XIII)

25       wherein R<sup>1</sup> is as defined above; by treatment with hydrogen bromide, typically in acetic acid at reflux.

The intermediates of formula XII and XIII above may be prepared by reacting a compound of formula III as defined above with the appropriate compound of formula XIV:




5

(XIV)

wherein M<sup>1</sup> is as defined above, and Y<sup>1</sup> represents amino or methoxy; in the presence of a transition metal catalyst; under conditions analogous to those described above for the reaction between compounds III and IV. In particular, the transition metal catalyst of use in the reaction between compounds III and XIV is suitably tetrakis(triphenylphosphine)-palladium(0), in which case the reaction is conveniently carried out at an elevated temperature in a solvent such as aqueous 1,2-dimethoxyethane, advantageously in the presence of sodium carbonate.

Where M<sup>1</sup> in the intermediates of formula V above represents -Sn(Alk)<sub>3</sub> and Alk is as defined above, this compound may be prepared by reacting a compound of formula III as defined above with a reagent of formula (Alk)<sub>3</sub>Sn-Hal, in which Hal represents a halogen atom, typically chloro. The reaction is conveniently effected by treating compound III with isopropylmagnesium chloride, typically in a solvent such as tetrahydrofuran, with subsequent addition of the stannyl reagent (Alk)<sub>3</sub>Sn-Hal.


Where L<sup>1</sup> in the intermediates of formula III above represents bromo, this compound may be prepared by bromination of the corresponding compound of formula XV:



(XV)

wherein R<sup>1</sup> is as defined above; typically by treatment with bromine in methanol, in the presence of sodium acetate and optionally also potassium  
5 bromide.

The intermediates of formula XV may be prepared by reacting chloroacetaldehyde or bromoacetaldehyde, or an acetal derivative thereof, e.g. the dimethyl or diethyl acetal thereof, with the requisite compound of  
10 formula XVI:




(XVI)

wherein R<sup>1</sup> is as defined above.

Where chloroacetaldehyde or bromoacetaldehyde is utilised as one  
15 of the reactants, the reaction is conveniently carried out by heating the reactants under basic conditions in a suitable solvent, e.g. sodium methoxide or sodium hydrogencarbonate in a lower alkanol such as methanol and/or ethanol at the reflux temperature of the solvent. Where  
an acetal derivative of chloroacetaldehyde or bromoacetaldehyde, e.g. the  
20 dimethyl or diethyl acetal thereof, is utilised as one of the reactants, the reaction is conveniently effected by heating the reactants under acidic conditions in a suitable solvent, e.g. aqueous hydrobromic acid in a lower alkanol such as methanol or ethanol, typically at the reflux temperature of the solvent.

In a still further procedure, the compounds according to the present invention may be prepared by a process which comprises reacting a compound of formula XVI as defined above with a compound of formula XVII:

5




(XVII)


wherein Y and Z are as defined above, and  $L^3$  represents a suitable leaving group; under conditions analogous to those described above for the reaction between chloroacetaldehyde or bromoacetaldehyde, or an acetal derivative thereof, and compound XVI.

The leaving group  $L^3$  is suitably a halogen atom, e.g. bromo.


The intermediates of formula XV may also be prepared by reacting a compound of formula XVIII or XIX with the compound of formula XX, or with an acid addition salt of the latter compound, e.g. the hemisulfate salt:



(XVIII)



(XIX)



(XX)


wherein  $R^1$  is as defined above, and  $Alk^1$  represents  $C_{1-6}$  alkyl.

Typical values of  $Alk^1$  include methyl and ethyl.


The reaction is conveniently effected by heating the reactants under basic conditions in a suitable solvent, e.g. a lower alkoxide such as sodium

methoxide or ethoxide in a lower alkanol such as methanol or ethanol, typically at the reflux temperature of the solvent.

In a yet further procedure, the compounds according to the present invention wherein  $R^1$  represents an aryl or heteroaryl moiety may be prepared by a process which comprises reacting a compound of formula XXI with a compound of formula XXII:



(XXI)



(XXII)

wherein Y, Z and  $M^1$  are as defined above,  $R^{1a}$  represents an aryl or heteroaryl moiety, and  $L^4$  represents a suitable leaving group; in the presence of a transition metal catalyst.

The leaving group  $L^4$  is typically a halogen atom, e.g. chloro.

The transition metal catalyst of use in the reaction between compounds XXI and XXII is suitably tetrakis(triphenylphosphine)-palladium(0), in which case the reaction is conveniently effected at an elevated temperature in a solvent such as *N,N*-dimethylacetamide, typically in the presence of potassium phosphate or in the presence of lithium chloride and copper(I) iodide. Alternatively, the transition metal catalyst may suitably be tris(dibenzylideneacetone)palladium(0), in which case the reaction is conveniently effected at an elevated temperature in a solvent such as 1,4-dioxane, typically in the presence of tri-*tert*-butylphosphine and cesium carbonate.

Where  $L^4$  in the compounds of formula XXII above represents a halogen atom, these compounds correspond to compounds of formula I as

defined above wherein  $R^1$  represents halogen, and they may therefore be prepared by any of the methods described above for the preparation of the compounds according to the invention.

5 The compound of formula XX above is commercially available from the Sigma-Aldrich Company Ltd., Dorset, United Kingdom.

Where they are not commercially available, the starting materials of formula VI, VIII, X, XIV, XVI, XVII, XVIII, XIX and XXI may be prepared by methods analogous to those described in the accompanying Examples, or by standard methods well known from the art.

10 It will be understood that any compound of formula I initially obtained from any of the above processes may, where appropriate, subsequently be elaborated into a further compound of formula I by techniques known from the art. For example, a compound of formula I wherein  $R^1$  represents  $-C(O-Alk^1)_2R^a$  initially obtained, wherein  $Alk^1$  is as  
15 defined above, may be converted into the corresponding compound of formula I wherein  $R^1$  represents  $-COR^a$  by hydrolysis with a mineral acid, typically aqueous hydrochloric acid. A compound wherein  $R^1$  represents formyl may be reduced with sodium triacetoxyborohydride to the corresponding compound wherein  $R^1$  represents hydroxymethyl. A  
20 compound of formula I wherein  $R^1$  represents hydroxymethyl may be oxidised to the corresponding compound of formula I wherein  $R^1$  represents formyl by treatment with manganese dioxide. The formyl derivative thereby obtained may be condensed with a hydroxylamine derivative of formula  $H_2N-OR^b$  to provide a compound of formula I wherein  
25  $R^1$  represents  $-CH=NOR^b$ . Furthermore, a compound of formula I wherein  $R^1$  represents  $-CH=NOH$  may be treated with triethylamine in the presence of 1,1'-carbonyldiimidazole to afford a corresponding compound of formula I wherein  $R^1$  represents cyano. Alternatively, the compound of formula I wherein  $R^1$  represents formyl may be reacted with a Grignard  
30 reagent of formula  $R^aMgBr$  to afford a compound of formula I wherein  $R^1$  represents  $-CH(OH)R^a$ , and this compound may in turn be oxidised using



manganese dioxide to the corresponding compound of formula I wherein  $R^1$  represents  $-\text{COR}^a$ . The latter compound may then be condensed with a hydroxylamine derivative of formula  $\text{H}_2\text{N}-\text{OR}^b$  to provide a compound of formula I wherein  $R^1$  represents  $-\text{CR}^a=\text{NOR}^b$ . A compound of formula I wherein  $R^1$  represents  $-\text{CH}(\text{OH})\text{R}^a$  may be converted into the corresponding compound of formula I wherein  $R^1$  represents  $-\text{CHFR}^a$  by treatment with (diethylamino)sulfur trifluoride (DAST). Similarly, a compound of formula I wherein  $R^1$  represents  $-\text{COR}^a$  may be converted into the corresponding compound of formula I wherein  $R^1$  represents  $-\text{CF}_2\text{R}^a$  by treatment with DAST. A compound of formula I wherein  $R^1$  represents amino may be converted into the corresponding compound of formula I wherein  $R^1$  represents chloro by diazotisation, using sodium nitrite, followed by treatment with copper(I) chloride. A compound of formula I wherein  $R^1$  represents  $-\text{COCH}_3$  may be treated with thioacetamide in the presence of pyridinium tribromide to furnish the corresponding compound of formula I wherein  $R^1$  represents 2-methylthiazol-5-yl. Moreover, a compound of formula I wherein  $R^1$  is formyl may be treated with (*p*-tolylsulfonyl)methyl isocyanide (TosMIC) in the presence of potassium carbonate to afford the corresponding compound of formula I wherein  $R^1$  represents oxazol-5-yl. A compound of formula I wherein  $R^1$  represents hydroxymethyl may be treated with carbon tetrabromide and triphenylphosphine to afford the corresponding compound of formula I wherein  $R^1$  represents bromomethyl, which may then be reacted (typically *in situ*) with the sodium salt of imidazole or 1*H*-[1,2,4]triazole to provide a compound of formula I wherein  $R^1$  represents imidazol-1-ylmethyl or [1,2,4]triazol-1-ylmethyl respectively; or with the sodium salt of 1*H*-[1,2,3]triazole to provide a mixture of compounds of formula I wherein  $R^1$  represents [1,2,3]triazol-1-ylmethyl and [1,2,3]triazol-2-ylmethyl; or with morpholine to provide a compound of formula I wherein  $R^1$  represents morpholin-4-ylmethyl.

Where a mixture of products is obtained from any of the processes described above for the preparation of compounds according to the invention, the desired product can be separated therefrom at an appropriate stage by conventional methods such as preparative HPLC; or  
5 column chromatography utilising, for example, silica and/or alumina in conjunction with an appropriate solvent system.

Where the above-described processes for the preparation of the compounds according to the invention give rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques  
10 such as preparative chromatography. The novel compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The novel compounds may, for example, be resolved into their component enantiomers by standard techniques such as preparative HPLC, or the formation of  
15 diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-*p*-toluoyl-d-tartaric acid and/or (+)-di-*p*-toluoyl-l-tartaric acid, followed by fractional crystallization and regeneration of the free base. The novel compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of  
20 the chiral auxiliary.

During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic*  
25 *Chemistry*, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The following Examples illustrate the preparation of compounds  
30 according to the invention.

The compounds in accordance with this invention potently inhibit the binding of [<sup>3</sup>H]-flumazenil to the benzodiazepine binding site of human GABA<sub>A</sub> receptors containing the  $\alpha 2$  and/or  $\alpha 3$  and/or  $\alpha 5$  subunit stably expressed in Ltk<sup>-</sup> cells.

5

### *Reagents*

- Phosphate buffered saline (PBS).
- Assay buffer: 10 mM KH<sub>2</sub>PO<sub>4</sub>, 100 mM KCl, pH 7.4 at room temperature.
- [<sup>3</sup>H]-Flumazenil (18 nM for  $\alpha 1\beta 3\gamma 2$  cells; 18 nM for  $\alpha 2\beta 3\gamma 2$  cells; 10 nM  
10 for  $\alpha 3\beta 3\gamma 2$  cells; 10 nM for  $\alpha 5\beta 3\gamma 2$  cells) in assay buffer.
- Flunitrazepam 100  $\mu$ M in assay buffer.
- Cells resuspended in assay buffer (1 tray to 10 ml).

### *Harvesting Cells*

- 15 Supernatant is removed from cells. PBS (approximately 20 ml) is added. The cells are scraped and placed in a 50 ml centrifuge tube. The procedure is repeated with a further 10 ml of PBS to ensure that most of the cells are removed. The cells are pelleted by centrifuging for 20 min at 3000 rpm in a benchtop centrifuge, and then frozen if desired. The pellets  
20 are resuspended in 10 ml of buffer per tray (25 cm x 25 cm) of cells.

### *Assay*

Can be carried out in deep 96-well plates or in tubes. Each tube contains:

- 25
- 300  $\mu$ l of assay buffer.
  - 50  $\mu$ l of [<sup>3</sup>H]-flumazenil (final concentration for  $\alpha 1\beta 3\gamma 2$ : 1.8 nM; for  $\alpha 2\beta 3\gamma 2$ : 1.8 nM; for  $\alpha 3\beta 3\gamma 2$ : 1.0 nM; for  $\alpha 5\beta 3\gamma 2$ : 1.0 nM).
  - 50  $\mu$ l of buffer or solvent carrier (e.g. 10% DMSO) if compounds are dissolved in 10% DMSO (total); test compound or flunitrazepam (to  
30 determine non-specific binding), 10  $\mu$ M final concentration.
  - 100  $\mu$ l of cells.

Assays are incubated for 1 hour at 40°C, then filtered using either a Tomtec or Brandel cell harvester onto GF/B filters followed by 3 x 3 ml washes with ice cold assay buffer. Filters are dried and counted by liquid scintillation counting. Expected values for total binding are 3000-4000 dpm for total counts and less than 200 dpm for non-specific binding if using liquid scintillation counting, or 1500-2000 dpm for total counts and less than 200 dpm for non-specific binding if counting with meltilex solid scintillant. Binding parameters are determined by non-linear least squares regression analysis, from which the inhibition constant  $K_i$  can be calculated for each test compound.

The compounds of the accompanying Examples were tested in the above assay, and all were found to possess a  $K_i$  value for displacement of [ $^3\text{H}$ ]-flumazenil from the  $\alpha 2$  and/or  $\alpha 3$  and/or  $\alpha 5$  subunit of the human GABA<sub>A</sub> receptor of 100 nM or less.

### EXAMPLE 1

#### 3'-(7-Methylimidazo[1,2-a]pyrimidin-3-yl)biphenyl-2-carbonitrile

A mixture of 2-bromobenzonitrile (9.1 g, 50 mmol), 3-aminobenzeneboronic acid monohydrate (11.6 g, 75 mmol) and tetrakis(triphenylphosphine)palladium(0) (1.73 g, 1.5 mmol) in 1,2-dimethoxyethane (50 ml) and 2M sodium carbonate solution (25 ml) was heated at 80°C for 20 h. After cooling to ambient temperature the reaction was partitioned between ethyl acetate (400 ml) and water (400 ml). The organics were washed with brine (400 ml), dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. Purification of the residue by chromatography on silica gel, eluting with isohexane on a gradient of ethyl acetate (0-25%) gave 3'-aminobiphenyl-2-carbonitrile as a colourless oil that solidified on standing to afford a white solid (9.5 g, 98%):  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 3.79 (2H, br), 6.75 (1H, ddd,  $J$  8, 3 and 1), 6.84 (1H, dd,  $J$  3 and 3), 6.92 (1H, dd,  $J$  8 and 3), 7.25 (1H, dd,  $J$  8 and 8), 7.40 (1H, ddd,  $J$

8, 8 and 1), 7.50 (1H, dd,  $J$  8 and 1), 7.62 (1H, ddd,  $J$  8, 8 and 1), 7.73 (1H, dd,  $J$  8 and 1).

A solution of 3'-aminobiphenyl-2-carbonitrile (10.9 g, 56 mmol) in 1,4-dioxane (30 ml) was treated with a solution of 25% aqueous sulfuric acid (150 ml). The resulting suspension was cooled to 0°C before being  
5 treated dropwise over 10 minutes with a solution of sodium nitrite (4.6 g, 67 mmol) in water (10 ml). After stirring at 0°C for 30 minutes the reaction was poured into hot (70°C) water (500 ml). On cooling to ambient temperature the product was extracted into ethyl acetate (500 ml), the  
10 organics were washed with water (300 ml), brine (300 ml) and dried over anhydrous sodium sulfate. Filtration and concentration *in vacuo* afforded 3'-hydroxybiphenyl-2-carbonitrile as a dark oil (7.1 g, 65%):  $\delta_H$  (400 MHz,  $CDCl_3$ ) 5.40 (1H, br), 6.92 (1H, ddd,  $J$  8, 3 and 1), 7.04 (1H, dd,  $J$  3 and 3), 7.11 (1H, ddd,  $J$  8, 3 and 1), 7.35 (1H, dd,  $J$  8 and 8), 7.44 (1H, ddd,  $J$  8, 8  
15 and 1), 7.51 (1H, dd,  $J$  8 and 1), 7.64 (1H, ddd,  $J$  8, 8 and 1), 7.75 (1H, dd,  $J$  8 and 1).

3'-Hydroxybiphenyl-2-carbonitrile (0.48 g, 2.47 mmol) and dry pyridine (0.98 g, 12.35 mmol) were dissolved in dichloromethane (7 ml) and cooled to 0°C before dropwise addition of trifluoromethanesulfonic  
20 anhydride (1.04 g, 3.70 mmol) over 5 min. The mixture was stirred at 0°C for 10 min and then at 25°C for 1 h. The solvent was evaporated *in vacuo* and the residue partitioned between ethyl acetate (200 ml) and water (150 ml). The organic layer was washed with brine (150 ml), dried over anhydrous sodium sulfate and evaporated to give a brown oil. Purification  
25 by silica gel chromatography eluting with isohexane on a gradient of ethyl acetate (0-30%) gave trifluoromethanesulfonic acid 2'-cyanobiphenyl-3-yl ester as a yellow oil (544 mg, 67%):  $\delta_H$  (400 MHz,  $CDCl_3$ ) 7.37 (1H, ddd,  $J$  8, 3 and 1), 7.39 (1H, dd,  $J$  3 and 3), 7.50-7.60 (2H, m), 7.61-7.65 (2H, m), 7.64 (1H, td,  $J$  8 and 1), 7.80 (1H, dd,  $J$  8 and 1).

30 Trifluoromethanesulfonic acid 2'-cyanobiphenyl-3-yl ester (0.55 g, 1.66 mmol), potassium acetate (0.49 g, 4.98 mmol) and

bis(pinacolato)diboron (0.55 g, 2.16 mmol) were dissolved in 1,4-dioxane (10 ml) and the mixture degassed with N<sub>2</sub> for 15 min. Dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane adduct (41 mg, 0.05 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (28 mg, 0.05 mmol) were then added and the mixture heated at 85°C for 18 h. The mixture was cooled to ambient temperature, filtered and the filter cake washed with diethyl ester. The filtrate was evaporated to dryness and partitioned between diethyl ether (25 ml) and 1N sodium hydroxide solution (25 ml). The aqueous layer was washed with more diethyl ether then made acidic (pH 6) with 4N hydrochloric acid. The resulting solid was collected by filtration and dried *in vacuo* to give 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as a cream-coloured solid (430 mg, 85%):  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 1.36 (12H, s), 7.43 (1H, ddd, *J* 8, 8 and 1.5), 7.48-7.57 (2H, m), 7.62 (2H, ddd, *J* 8, 8 and 1.5), 7.68-7.71 (1H, m), 7.74-7.77 (1H, m), 7.87-7.90 (1H, m), 7.92-7.94 (1H, m).

Sodium methoxide (1.62 g, 30 mmol) was added to a stirred solution of 2-aminoimidazole hemisulfate (2.64 g, 20 mmol) and 1,1-dimethoxy-3-butanone (2 ml) in ethanol (25 ml). The mixture was heated under reflux for 8 h, allowed to cool to room temperature then pre-adsorbed directly onto silica. Purification by silica gel chromatography eluting with dichloromethane and 1% conc. ammonia on a gradient of methanol (1-4%) gave a 95:5 mixture of 7-methylimidazo[1,2-*a*]pyrimidine and 5-methylimidazo[1,2-*a*]pyrimidine respectively (1.68 g, 64%) as a white crystalline solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>, 7-methyl isomer) 2.64 (3H, s), 6.74 (1H, d, *J* 7), 7.45 (1H, d, *J* 1), 7.73 (1H, d, *J* 1), 8.29 (1H, d, *J* 7).

7-Methylimidazo[1,2-*a*]pyrimidine (100 mg, 0.75 mmol) and sodium acetate (74 mg, 0.90 mmol) were dissolved in methanol (2 ml) which had been saturated with potassium bromide and this mixture was cooled to -10°C before dropwise addition of bromine (132 mg, 0.83 mmol) over 5 min. On complete addition the mixture was quenched by addition of 1M sodium sulfite solution (2 ml) and the solvent removed *in vacuo*. The residue was

treated with water (15 ml) and saturated sodium hydrogencarbonate solution (15 ml) and extracted with ethyl acetate (3 x 40 ml). The organics were combined then washed with brine (40 ml), dried over anhydrous sodium sulfate and evaporated to give an off white solid. This solid was  
5 purified by silica gel chromatography eluting with dichloromethane and 1% conc. ammonia on a gradient of methanol (1-2%) to give 3-bromo-7-methylimidazo[1,2-*a*]pyrimidine (100 mg, 63%) as a white crystalline solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 2.67 (3H, s), 6.87 (1H, d, *J* 7), 7.71 (1H, s), 8.27 (1H, d, *J* 7).

10 3-Bromo-7-methylimidazo[1,2-*a*]pyrimidine (100 mg, 0.47 mmol), potassium phosphate (200 mg, 0.94 mmol) and 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile (215 mg, 0.70 mmol) were dissolved in *N,N*-dimethylacetamide (2 ml) and the mixture degassed with  $N_2$  for 15 min. Tetrakis(triphenylphosphine)palladium(0) (27 mg, 0.002  
15 mmol) was added and the mixture heated at 65°C for 24 h. The mixture was allowed to cool to room temperature, diluted with water (20 ml) and saturated sodium hydrogencarbonate solution (20 ml) then extracted with ethyl acetate (2 x 75 ml). The combined organic fractions were washed  
20 with brine (40 ml), dried over anhydrous sodium sulfate and evaporated to give a black oil. The oil was purified by silica gel chromatography eluting with dichloromethane and 1% conc. ammonia on a gradient of methanol (1-2%). The solid obtained was triturated with diethyl ether to furnish 3'-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile (120 mg, 82%)  
25 as a white powder:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 2.67 (3H, s), 6.83 (1H, d, *J* 7), 7.49 (1H, td, *J* 8 and 1), 7.55-7.75 (6H, m), 7.82 (1H, dd, *J* 8 and 1), 7.85 (1H, s), 8.82 (1H, d, *J* 7).

## EXAMPLE 2

30 3'-(Imidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile

A solution of 2-aminopyrimidine (0.5 g, 5.26 mmol), bromoacetaldehyde diethyl acetal (2.07 g, 10.5 mmol) and 48% aqueous hydrobromic acid (0.5 ml) in ethanol (5 ml) was heated at reflux for 18 h. The reaction was cooled and pre-adsorbed directly onto silica gel.

5 Purification by flash chromatography eluting with dichloromethane (containing 1% conc. ammonia) on a gradient of methanol (1-3%) gave a solid which was triturated with 5% diethyl ether in isohexane to afford imidazo[1,2- $\alpha$ ]pyrimidine (0.51 g, 82%) as a tan solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 6.92 (1H, dd,  $J$  7 and 4), 7.59 (1H, d,  $J$  1), 7.84 (1H, d,  $J$  1), 8.49 (1H, dd,  $J$  7 and 2), 8.58 (1H, dd,  $J$  7 and 2).

Imidazo[1,2- $\alpha$ ]pyrimidine (0.20 g, 1.68 mmol) was brominated as described in Example 1 to give 3-bromoimidazo[1,2- $\alpha$ ]pyrimidine (0.29 g, 87%) as a white solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 7.02 (1H, dd,  $J$  7 and 4), 7.83 (1H, s), 8.43 (1H, dd,  $J$  7 and 2), 8.59 (1H, dd,  $J$  7 and 2).

15 3-Bromoimidazo[1,2- $\alpha$ ]pyrimidine (0.29 g, 1.68 mmol) was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-(imidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (260 mg, 60%) as an off-white powder:  $\delta_H$  (360 MHz,  $CDCl_3$ ) 6.97 (1H, dd,  $J$  7 and 4), 7.51 (1H, td,  $J$  1 and 8), 7.59 (1H, dd,  $J$  7 and 1), 7.62-7.73 (4H, m), 7.77-7.78 (1H, m), 7.82 (1H, dd,  $J$  8 and 1), 7.96 (1H, s), 8.60 (1H, dd,  $J$  7 and 2), 8.98 (1H, d,  $J$  7 and 2);  $m/z$  ( $ES^+$ ) 297 ( $M^++H$ ).

### EXAMPLE 3

#### 3'-(7-Trifluoromethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile

25 A mixture of 2-amino-4-(trifluoromethyl)pyrimidine (prepared according to Zanatta *et al.* in *J. Heterocyclic Chem.*, 1997, 34(2), 509-513) (500 mg, 3.1 mmol) and bromoacetaldehyde diethyl acetal (1.38 ml, 9.2 mmol) in ethanol (10 ml) was treated with hydrobromic acid (0.5 ml of a 48% aqueous solution) and then heated at 70°C for 12 h. The reaction was



cooled to ambient temperature then pre-adsorbed onto silica. Purification by chromatography on silica eluting with dichloromethane (containing 1% conc. ammonia) on a gradient of methanol (1-5%) afforded 7-

trifluoromethylimidazo[1,2-*a*]pyrimidine (500 mg, 87%) as a cream-

5 coloured solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 7.22 (1H, d, *J* 7), 7.74 (1H, d, *J* 1), 8.03 (1H, d, *J* 1), 8.67 (1H, d, *J* 7).

7-Trifluoromethylimidazo[1,2-*a*]pyrimidine (0.20 g, 1.07 mmol) was brominated as described in Example 1 to give 3-bromo-7-

10 trifluoromethylimidazo[1,2-*a*]pyrimidine (0.28 g, 98%) as a white solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 7.35 (1H, d, *J* 7), 8.02 (1H, s), 8.62 (1H, d, *J* 7).

3-Bromo-7-trifluoromethylimidazo[1,2-*a*]pyrimidine (0.28 g, 1.04 mmol) was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-(7-trifluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile (165  
15 mg, 44%) as a yellow powder:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 7.29 (1H, d, *J* 7), 7.53 (1H, td, *J* 8 and 1), 7.59-7.68 (3H, m), 7.70-7.75 (2H, m), 7.80-7.85 (2H, m), 8.14 (1H, s), 9.20 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 365 (M<sup>+</sup>+H).

#### EXAMPLE 4

20 3'-[7-(1,1-Dimethoxyethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile

Boron trifluoride etherate (17.03 g, 120.0 mmol) was added dropwise over 15 min to a cooled (-40°C) solution of triethyl orthoformate  
25 (14.82 g, 100.0 mmol) in dichloromethane (50 ml). Stirring was continued for 10 min then the solution was transferred to an ice-water bath and stirred at 0°C for 20 min. The mixture was cooled to -78°C, and 3,3-dimethoxy-2-butanone (6.61 g, 50.0 mmol) added followed by dropwise addition of *N,N*-diisopropylethylamine (19.39 g, 150.0 mmol) over 15 min.  
30 Stirring was continued for 1 h then the solution was poured onto a vigorously stirred mixture of saturated sodium hydrogencarbonate

solution (500 ml) and dichloromethane (200 ml). The organic phase was separated, washed with ice-cold 1M sulfuric acid solution (2 x 500 ml) and ice-cold water (2 x 500 ml), dried over anhydrous sodium sulfate solution and evaporated to give 1,1-diethoxy-4,4-dimethoxypentan-3-one (11.72 g,  
5 100%) as an orange oil.

1,1-Diethoxy-4,4-dimethoxypentan-3-one was condensed with 2-aminoimidazole hemisulfate as described in Example 1 to give 7-(1,1-dimethoxyethyl)imidazo[1,2-*a*]pyrimidine (6.61 g, 64%) as a white solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 1.70 (3H, s), 3.28 (6H, s), 7.30 (1H, d, *J* 7), 7.55 (1H, d, *J* 1), 7.84 (1H, d, *J* 1), 8.43 (1H, d, *J* 7).  
10

7-(1,1-Dimethoxyethyl)imidazo[1,2-*a*]pyrimidine (207 mg, 1.00 mmol) was brominated as described in Example 1 to give 3-bromo-7-(1,1-dimethoxyethyl)imidazo[1,2-*a*]pyrimidine (197 mg, 69%) as a white solid:  $\delta_H$  (360 MHz,  $CDCl_3$ ) 1.70 (3H, s), 3.28 (6H, s), 7.43 (1H, d, *J* 7), 7.82 (1H, s), 8.39 (1H, d, *J* 7).  
15

3-Bromo-7-(1,1-dimethoxyethyl)imidazo[1,2-*a*]pyrimidine (197 mg, 0.50 mmol) was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-[7-(1,1-dimethoxyethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile (200  
20 mg, 76%) as a white powder:  $\delta_H$  (360 MHz,  $CDCl_3$ ) 1.72 (3H, s), 3.30 (6H, s), 7.39 (1H, d, *J* 7), 7.49-7.73 (6H, m), 7.78-7.84 (2H, m), 7.97 (1H, s), 8.95 (1H, d, *J* 7); *m/z* ( $ES^+$ ) 385 ( $M^+ + H$ ).

### EXAMPLE 5

25

#### 3'-(7-Acetylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile

A solution of 3'-[7-(1,1-dimethoxyethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile (100 mg, 0.26 mmol) in 2.5N hydrochloric acid (4 ml) was heated at 50°C for 15 h. The mixture was cooled, layered with ethyl acetate (5 ml) and made basic (pH 8) with portionwise addition of  
30 solid sodium hydrogencarbonate over 15 min. The mixture was diluted

with water (5 ml) and extracted with dichloromethane (5 x 30 ml). The combined organics were dried over anhydrous sodium sulfate, filtered, evaporated to dryness and triturated with diethyl ether to give 3'-(7-acetylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile (62 mg, 70%) as a yellow solid:  $\delta_H$  (360 MHz,  $CDCl_3$ ) 2.84 (3H, s), 7.26 (1H, d, *J* 7), 7.52 (1H, td, *J* 8 and 1), 7.58-7.72 (5H, m), 7.80-7.84 (2H, m), 8.16 (1H, s), 9.03 (1H, d, *J* 7); *m/z* ( $ES^+$ ) 339 ( $M^+ + H$ ).

### EXAMPLE 6

#### 3'-(7-Isopropylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile

3-Methylbutan-2-one was converted to 1,1-diethoxy-4-methylpentan-3-one as described in Example 4 and condensed with 2-aminoimidazole hemisulfate as described in Example 1 to give 7-

isopropylimidazo[1,2-*a*]pyrimidine as an orange solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 1.36 (6H, d, *J* 7), 3.12 (1H, septet, *J* 7), 6.78 (1H, d, *J* 7), 7.45 (1H, d, *J* 1), 7.72 (1H, d, *J* 1), 8.33 (1H, d, *J* 7).

7-Isopropylimidazo[1,2-*a*]pyrimidine was brominated as described in Example 1 to give 3-bromo-7-isopropylimidazo[1,2-*a*]pyrimidine as a cream-coloured solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 1.37 (6H, d, *J* 7), 3.16 (1H, septet, *J* 7), 6.91 (1H, d, *J* 7), 7.71 (1H, s), 8.30 (1H, d, *J* 7).

3-Bromo-7-isopropylimidazo[1,2-*a*]pyrimidine was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-(7-isopropylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile as a white powder: Found C, 76.71; H, 5.17; N 16.24.  $C_{22}H_{18}N_4 \cdot 0.3(H_2O)$  requires C, 76.59; H, 5.43; N, 16.24;  $\delta_H$  (400 MHz,  $CDCl_3$ ) 1.38 (6H, d, *J* 7), 3.16 (1H, septet, *J* 7), 6.88 (1H, d, *J* 7), 7.48-7.76 (7H, m), 7.82 (1H, dd, *J* 8 and 1), 7.85 (1H, s), 8.85 (1H, d, *J* 7); *m/z* ( $ES^+$ ) 339 ( $M^+ + H$ ).

### EXAMPLE 7

#### 3'-(7-Cyclopropylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile

Cyclopropyl methyl ketone was converted to 1-cyclopropyl-3,3-diethoxypropan-1-one as described in Example 4 and condensed with 2-aminoimidazole hemisulfate as described in Example 1 to give 7-cyclopropylimidazo[1,2- $\alpha$ ]pyrimidine as an orange solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 1.06-1.11 (2H, m), 1.26-1.31 (2H, m), 2.03-2.08 (1H, m), 6.74 (1H, d,  $J$  7), 7.40 (1H, d,  $J$  1), 7.65 (1H, d,  $J$  1), 8.24 (1H, d,  $J$  7).

7-Cyclopropylimidazo[1,2- $\alpha$ ]pyrimidine was brominated as described in Example 1 to give 3-bromo-7-cyclopropylimidazo[1,2- $\alpha$ ]pyrimidine as a tan solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 1.10-1.15 (2H, m), 1.29-1.32 (2H, m), 2.04-2.12 (1H, m), 6.88 (1H, d,  $J$  7), 7.63 (1H, s), 8.20 (1H, d,  $J$  7).

3-Bromo-7-cyclopropylimidazo[1,2- $\alpha$ ]pyrimidine was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-(7-cyclopropylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile. Hydrochloride salt, white powder (from acetonitrile):  $\delta_H$  (400 MHz, DMSO) 1.21-1.25 (2H, m), 1.31-1.36 (2H, m), 2.46-2.52 (1H, m), 7.58 (1H, d,  $J$  7), 7.65 (1H, td,  $J$  7 and 1), 7.77-7.88 (5H, m), 7.93-7.94 (1H, m), 8.01 (1H, dd,  $J$  7 and 1), 8.37 (1H, s), 9.16 (1H, d,  $J$  7);  $m/z$  (ES<sup>+</sup>) 337 (M<sup>+</sup>+H).

### EXAMPLE 8

#### 3'-(7-*tert*-Butylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile

3,3-Dimethylbutan-2-one was converted to 1,1-diethoxy-4,4-dimethylpentan-3-one as described in Example 4 and condensed with 2-aminoimidazole hemisulfate as described in Example 1 to give 7-*tert*-butylimidazo[1,2- $\alpha$ ]pyrimidine as a pale-orange solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 1.42 (9H, s), 6.96 (1H, d,  $J$  7), 7.45 (1H, d,  $J$  1), 7.72 (1H, d,  $J$  1), 8.33 (1H, d,  $J$  7).

7-*tert*-Butylimidazo[1,2-*a*]pyrimidine was brominated as described in Example 1 to give 3-bromo-7-*tert*-butylimidazo[1,2-*a*]pyrimidine as an off-white solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 1.43 (9H, s), 7.09 (1H, d, *J* 7), 7.71 (1H, s), 8.30 (1H, d, *J* 7).

5        3-Bromo-7-*tert*-butylimidazo[1,2-*a*]pyrimidine was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-(7-*tert*-butylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile as an off-white powder: Found C, 77.17; H, 5.79; N, 15.78.  $C_{23}H_{20}N_4 \cdot 0.3(H_2O)$  requires C, 77.21; H, 5.80; N, 15.66;  $\delta_H$  (400  
10 MHz,  $CDCl_3$ ) 1.44 (9H, s), 7.05 (1H, d, *J* 7), 7.48-7.76 (7H, m), 7.83 (1H, dd, *J* 8 and 1), 7.86 (1H, s), 8.88 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 353 ( $M^+ + H$ ).

### EXAMPLE 9

#### 15    3'-(7-Cyclobutylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile

Cyclobutyl methyl ketone was converted to 1-cyclobutyl-3,3-diethoxypropan-1-one as described in Example 4 and condensed with 2-aminoimidazole hemisulfate as described in Example 1 to give 7-cyclobutylimidazo[1,2-*a*]pyrimidine as an orange solid:  $\delta_H$  (400 MHz,  
20  $CDCl_3$ ) 1.90-2.55 (6H, m), 3.71 (1H, quintet, *J* 8), 6.73 (1H, d, *J* 7), 7.45 (1H, d, *J* 1), 7.72 (1H, d, *J* 1), 8.30 (1H, dd, *J* 7 and 3).

7-Cyclobutylimidazo[1,2-*a*]pyrimidine was brominated as described in Example 1 to give 3-bromo-7-cyclobutylimidazo[1,2-*a*]pyrimidine as a cream-coloured solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 1.90-2.55 (6H, m), 3.74 (1H,  
25 quintet, *J* 8), 6.85 (1H, d, *J* 7), 7.71 (1H, s), 8.27 (1H, d, *J* 7).

3-Bromo-7-cyclobutylimidazo[1,2-*a*]pyrimidine was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-(7-cyclobutylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile as a white powder:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 1.90-  
30 2.55 (6H, m), 3.77 (1H, quintet, *J* 8), 6.83 (1H, d, *J* 7), 7.48-7.75 (7H, m),

7.82 (1H, dd,  $J$  8 and 1), 7.86 (1H, s), 8.85 (1H, d,  $J$  7);  $m/z$  (ES<sup>+</sup>) 351 (M<sup>+</sup>+H).

### EXAMPLE 10

5

#### 3'-(7-Methoxyimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile

2-Amino-4-chloropyrimidine (1.00 g, 7.72 mmol) and sodium methoxide (1.25 g, 23.2 mmol) were suspended in methanol (20 ml) and heated under reflux for 2 h then cooled to ambient temperature.

10 Chloroacetaldehyde (2.4 ml of a 45% w/w solution in water, 15.44 mmol) was added and the mixture heated under reflux for 22 h, allowed to cool to ambient temperature then pre-adsorbed directly onto silica. Purification by chromatography on silica gel eluting with dichloromethane (containing 1% conc. ammonia) on a gradient of methanol (1-4%) gave 7-  
15 methoxyimidazo[1,2- $\alpha$ ]pyrimidine (0.77 g, 67%) as a white crystalline solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 4.04 (3H, s), 6.41 (1H, d,  $J$  7), 7.31 (1H, d,  $J$  1), 7.50 (1H, d,  $J$  1), 8.17 (1H, d,  $J$  7).

7-Methoxyimidazo[1,2- $\alpha$ ]pyrimidine (0.66 g, 4.43 mmol) was brominated as described in Example 1 to give 3-bromo-7-  
20 methoxyimidazo[1,2- $\alpha$ ]pyrimidine (0.48 g, 48%) as an off-white solid:  $\delta_H$  (360 MHz, CDCl<sub>3</sub>) 4.06 (3H, s), 6.53 (1H, d,  $J$  7), 7.48 (1H, s), 8.14 (1H, d,  $J$  7).

3-Bromo-7-methoxyimidazo[1,2- $\alpha$ ]pyrimidine (0.48 g, 2.12 mmol) was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-  
25 2-carbonitrile as described in Example 1 to give 3'-(7-methoxyimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (458 mg, 66%) as a white powder:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 4.08 (3H, s), 6.49 (1H, d,  $J$  7), 7.50 (1H, td,  $J$  8 and 1), 7.51-7.71 (8H, m), 7.81 (1H, dd,  $J$  8 and 1), 8.66 (1H, d,  $J$  7);  $m/z$  (ES<sup>+</sup>) 327 (M<sup>+</sup>+H).

30

EXAMPLE 113'-(7-Hydroxymethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile

Pyruvic aldehyde dimethyl acetal (8.43 g, 71.4 mmol) and *N,N*-  
5 dimethylformamide dimethyl acetal (8.51 g, 71.4 mmol) were heated at  
100°C for 18 h. The mixture was concentrated to give a brown oil and was  
then added dropwise over 10 min to a warm (60°C) suspension of 2-  
aminoimidazole hemisulfate (9.43 g, 71.4 mmol) in water (50 ml). The  
mixture was heated at 50°C for 36 h, cooled to ambient temperature and  
10 then pre-adsorbed directly onto silica. Purification by chromatography on  
silica gel eluting with dichloromethane (containing 1% conc. ammonia) on  
a gradient of methanol (1-2%) gave a 3:1 mixture of 7-  
dimethoxymethylimidazo[1,2- $\alpha$ ]pyrimidine and 5-  
dimethoxymethylimidazo[1,2- $\alpha$ ]pyrimidine respectively. Crystallisation  
15 from toluene gave 7-dimethoxymethylimidazo[1,2- $\alpha$ ]pyrimidine (2.20 g,  
16%) as a brown crystalline solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 3.50 (6H, s), 5.26  
(1H, s), 7.15 (1H, d, *J* 7), 7.56 (1H, d, *J* 1), 7.84 (1H, d, *J* 1), 8.47 (1H, d, *J*  
7).

7-Dimethoxymethylimidazo[1,2- $\alpha$ ]pyrimidine (1.00 g, 5.18 mmol)  
20 was dissolved in 3N hydrochloric acid and heated at 48°C for 14 h. The  
solution was layered with ethyl acetate (30 ml) and solid sodium  
hydrogencarbonate (1.06 g, 12.6 mmol) was added in portions over 5 min.  
The mixture was diluted with water (6 ml) and extracted with  
dichloromethane (5 x 50 ml). The combined organics were dried over  
25 anhydrous sodium sulfate, filtered and evaporated to give imidazo[1,2- $\alpha$ ]  
pyrimidine-7-carbaldehyde (749 mg, 99%) as a yellow solid:  $\delta_H$  (360 MHz,  
CDCl<sub>3</sub>) 7.53 (1H, d, *J* 7), 7.77 (1H, d, *J* 1), 8.10 (1H, d, *J* 1), 8.60 (1H, d, *J*  
7), 10.05 (1H, s).

Sodium triacetoxyborohydride (21.5 g, 102 mmol) was added  
30 portionwise over 20 min to a stirred solution of imidazo[1,2- $\alpha$ ]pyrimidine-  
7-carbaldehyde (5.00 g, 34.0 mmol) in methanol (100 ml) and the solution

left to stir at ambient temperature for 18 h. The solvent was evaporated, the residue redissolved in methanol (150 ml) and pre-adsorbed onto silica. Purification by chromatography on silica gel eluting with dichloromethane (containing 1% conc. ammonia) on a gradient of methanol (1-10%) gave  
5 imidazo[1,2- $\alpha$ ]pyrimidin-7-ylmethanol (5.06 g, 99%) as a white solid:  $\delta_H$  (360 MHz, DMSO) 4.57 (2H, d,  $J$  6), 5.62 (1H, t,  $J$  6), 7.13 (1H, d,  $J$  7), 7.64 (1H, d,  $J$  1), 7.86 (1H, d,  $J$  1), 8.94 (1H, d,  $J$  7).

Imidazo[1,2- $\alpha$ ]pyrimidin-7-ylmethanol (1.70 g, 11.4 mmol) was brominated as described in Example 1 to give 3-bromoimidazo[1,2- $\alpha$ ]pyrimidin-7-ylmethanol (912 mg, 35%) as a white solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 2.95 (1H, br), 4.88 (2H, s), 7.03 (1H, d,  $J$  7), 7.73 (1H, s), 8.37 (1H, d,  $J$  7).  
10

3-Bromoimidazo[1,2- $\alpha$ ]pyrimidin-7-ylmethanol (912 mg, 4.00 mmol) was dissolved in dichloromethane (10 ml) and treated with imidazole (0.70 g, 10.3 mmol) and *tert*-butyldimethylsilyl chloride (0.77 g, 5.1 mmol) and the mixture left to stir at ambient temperature for 1 h. The reaction was diluted with dichloromethane (75 ml) and washed with 0.01N hydrochloric acid (2 x 50 ml). The organic phase was washed with saturated sodium hydrogencarbonate solution (50 ml), water (50 ml) and brine (50 ml), dried  
15 over anhydrous magnesium sulfate, filtered and evaporated to give crude 3-bromo-7-(*tert*-butyldimethylsilyloxymethyl)imidazo[1,2- $\alpha$ ]pyrimidine (1.37 g, 100%) as a yellow solid.  
20

3-Bromo-7-(*tert*-butyldimethylsilyloxymethyl)imidazo[1,2- $\alpha$ ]pyrimidine (1.37 g, 4.00 mmol) was coupled with 3'-(4,4,5,5-tetramethyl-  
25 [1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-[7-(*tert*-butyldimethylsilyloxymethyl)imidazo[1,2- $\alpha$ ]pyrimidin-3-yl]biphenyl-2-carbonitrile as a yellow solid. This solid was suspended in methanol (25 ml) and treated with conc. hydrochloric acid (1 ml) and left to stir at ambient temperature for 5 min. The solution was loaded onto a  
30 cartridge of strong cation-exchange resin, eluting with methanol then with 10% conc. ammonia in methanol. The basic fractions were concentrated *in*



*vacuo* and the residue purified further by chromatography on silica gel. Elution with dichloromethane (containing 1% conc. ammonia) on a gradient of methanol (2-5%) afforded 3'-(7-hydroxymethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (587 mg, 45%) as a yellow powder:  
5  $\delta_H$  (400 MHz,  $CDCl_3$ ) 1.90 (1H, s), 4.88 (2H, s), 6.91 (1H, d,  $J$  7), 7.51 (1H, td,  $J$  8 and 1), 7.57-7.73 (5H, m), 7.77 (1H, dd,  $J$  1 and 1), 7.82 (1H, dd,  $J$  8 and 1), 7.89 (1H, s), 8.94 (1H, d,  $J$  7);  $m/z$  ( $ES^+$ ) 327 ( $M^++H$ ).

### EXAMPLE 12

#### 3'-(7-Fluoromethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile

To a cooled ( $-78^\circ C$ ) suspension of 3'-(7-hydroxymethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (75 mg, 0.23 mmol) in dichloromethane (20 ml) was added (diethylamino)sulfur trifluoride (39  
15 mg, 0.24 mmol) dropwise over 5 min. The mixture was stirred at  $-78^\circ C$  for 20 min then allowed to warm to  $-40^\circ C$  for 5 min before quenching the reaction with a pre-cooled ( $-40^\circ C$ ) solution of acetic acid (0.5 ml) in dichloromethane (5 ml). The mixture was warmed to ambient temperature, made basic with saturated sodium hydrogencarbonate  
20 solution (30 ml) and extracted with dichloromethane (2 x 50 ml). The combined organics were washed with brine (30 ml), dried over anhydrous sodium sulfate and evaporated to dryness. Purification of this material by preparative thin-layer chromatography eluting with  
dichloromethane/methanol/conc. ammonia (96:4:0.4) gave a solid which  
25 was triturated with diethyl ether to give 3'-(7-fluoromethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (8 mg, 11%) as a yellow powder:  
 $\delta_H$  (360 MHz,  $CDCl_3$ ) 5.57 (2H, d,  $J$  47), 7.20 (1H, dd,  $J$  7 and 2), 7.52 (1H, dt,  $J$  8 and 1), 7.58-7.74 (5H, m), 7.78-7.79 (1H, m), 7.83 (1H, d,  $J$  8), 7.95 (1H, s), 9.03 (1H, dd,  $J$  7 and 1);  $m/z$  ( $ES^+$ ) 329 ( $M^++H$ ).

**EXAMPLE 13****3'-(7-Formylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile**

3'-(7-Hydroxymethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (20 mg, 0.061 mmol) and manganese(IV) oxide (53 mg, 0.61 mmol) were suspended in 1,2-dichloroethane (2 ml) and heated at 50°C for 18 h. The mixture was filtered through Celite and the filtrate concentrated *in vacuo*. Purification of the residue by preparative thin-layer chromatography eluting with dichloromethane/methanol/conc. ammonia (96:3:0.3) gave a solid which was triturated with diethyl ether to give 3'-(7-formylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (10 mg, 50%) as a yellow powder:  $\delta_H$  (360 MHz, DMSO) 7.48 (1H, d,  $J$  7), 7.64 (1H, td,  $J$  8 and 1), 7.70-7.92 (5H, m), 7.95-8.05 (2H, m), 8.42 (1H, s), 9.32 (1H, d,  $J$  7), 9.96 (1H, s);  $m/z$  (ES<sup>+</sup>) 325 (M<sup>+</sup>+H).

**EXAMPLE 14****3'-(7-Hydroxyiminomethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile**

3'-(7-Formylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (100 mg, 0.31 mmol) and 50% aqueous hydroxylamine (127 mg, 1.54 mmol) were dissolved in ethanol (2 ml) and the mixture heated at 40°C for 1 h. Purification of the reaction mixture by preparative thin-layer chromatography eluting with dichloromethane/methanol/conc. ammonia (96:4:0.4) gave a solid which was triturated with diethyl ether to give 3'-(7-hydroxyiminomethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (70 mg, 67%) as a white powder:  $\delta_H$  (400 MHz, DMSO) 7.42 (1H, d,  $J$  7), 7.61-7.80 (5H, m), 7.82-7.86 (2H, m), 7.94 (1H, dd,  $J$  1 and 1), 8.00 (1H, dd,  $J$  8 and 1), 8.10 (1H, d,  $J$  8), 9.09 (1H, d,  $J$  7), 12.20 (1H, s);  $m/z$  (ES<sup>+</sup>) 340 (M<sup>+</sup>+H).

**EXAMPLE 15****3-(2'-Cyanobiphenyl-3-yl)imidazo[1,2- $\alpha$ ]pyrimidine-7-carbonitrile**

3'-(7-Hydroxyiminomethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-  
5 carbonitrile (60 mg, 0.177 mmol) and 1,1'-carbonyldiimidazole (143 mg,  
0.88 mmol) were suspended in dichloromethane (2 ml) and triethylamine  
(89 mg, 0.88 mmol) was added dropwise over 5 min. Stirring was  
continued for 15 min then the mixture was heated under reflux for 90 min.  
Purification of the reaction mixture by preparative thin-layer  
10 chromatography eluting with dichloromethane/methanol/conc. ammonia  
(98:2:0.2) gave a solid which was triturated with diethyl ether to give 3-(2'-  
cyanobiphenyl-3-yl)imidazo[1,2- $\alpha$ ]pyrimidine-7-carbonitrile (56 mg, 97%)  
as a white powder:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 7.26 (1H, d,  $J$  7), 7.53 (1H, td,  $J$  8  
and 1), 7.59-7.67 (3H, m), 7.70-7.76 (2H, m), 7.81-7.85 (2H, m), 8.24 (1H,  
15 s), 9.16 (1H, d,  $J$  7);  $m/z$  (ES<sup>+</sup>) 322 (M<sup>++</sup>H).

**EXAMPLE 16****3-(2'-Methoxybiphenyl-3-yl)-7-methylimidazo[1,2- $\alpha$ ]pyrimidine**

20 A mixture of 3-bromo-7-methylimidazo[1,2- $\alpha$ ]pyrimidine (5.3 g, 25  
mmol), 3-methoxyphenylboronic acid (4.94 g, 32.5 mmol) and  
tetrakis(triphenylphosphine)palladium(0) (870 mg, 0.8 mmol) in 1,2-  
dimethoxyethane (40 ml) was treated with sodium carbonate (18.8 ml of a  
2M aqueous solution) then heated at 85°C for 16 h. The reaction mixture  
25 was allowed to cool to ambient temperature then partitioned between  
ethyl acetate and water. The organics were washed with 10% sodium  
carbonate solution, water, brine, dried over anhydrous magnesium sulfate,  
filtered then pre-adsorbed onto silica. Purification by chromatography on  
silica gel eluting with dichloromethane:conc. ammonia (99.5:0.5) on a  
30 gradient of methanol (1-3%) afforded 3-(3-methoxyphenyl)-7-  
methylimidazo[1,2- $\alpha$ ]pyrimidine as a cream-coloured solid (5.3 g, 89%):  $\delta_H$

(400 MHz, CDCl<sub>3</sub>) 2.66 (3H, s), 3.87 (3H, s), 6.76 (1H, d, *J* 7), 6.95-6.98 (1H, m), 7.03-7.05 (1H, m), 7.09-7.11 (1H, m), 7.44 (1H, t, *J* 8), 7.80 (1H, s), 8.52 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 240 (M<sup>+</sup>+H).

5 A mixture of 3-(3-methoxyphenyl)-7-methylimidazo[1,2-  
α]pyrimidine (5.02 g, 21 mmol) in hydrogen bromide (25 ml of a 30 wt. %  
solution in acetic acid) was heated at 130°C for 8 h. A further portion of  
hydrogen bromide solution was added (25 ml) and heating continued for 16  
h. The reaction was cooled to ambient temperature and diluted with ice-  
cold water (200 ml). The aqueous phase was made neutral with 4N  
10 sodium hydroxide then extracted with dichloromethane (2 x 250 ml). The  
organics were combined, washed with brine, dried over anhydrous  
magnesium sulfate, filtered and evaporated to dryness. The residue was  
trituated with ether to afford 3-(7-methylimidazo[1,2-α]pyrimidin-3-yl)-  
phenol as a light-grey solid (4.4 g, 93%): *m/z* (ES<sup>+</sup>) 226 (M<sup>+</sup>+H).

15 A cooled (-10°C) suspension of 3-(7-methylimidazo[1,2-α]pyrimidin-  
3-yl)phenol (4.28 g, 19 mmol) in dichloromethane (75 ml) was treated with  
pyridine (2.46 ml, 30 mmol) then with trifluoromethanesulfonic anhydride  
(4.05 ml, 25 mmol) added dropwise over 15 min. The resulting dark  
solution was stirred at -10°C for 30 min then warmed to ambient  
20 temperature. The reaction was diluted with dichloromethane (200 ml)  
then extracted with 0.1N hydrochloric acid (2 x 200ml), water, brine, dried  
over anhydrous sodium sulfate, filtered and evaporated to dryness to  
afford crude trifluoromethanesulfonic acid 3-(7-methylimidazo[1,2-  
α]pyrimidin-3-yl)phenyl ester as a dark oil: *m/z* (ES<sup>+</sup>) 316 (M<sup>+</sup>+H). This oil  
25 was dissolved in *N,N*-dimethylacetamide (50 ml) and degassed with  
nitrogen for 30 min before adding potassium phosphate (8.1 g, 38 mmol),  
tetrakis(triphenylphosphine)palladium(0) (660 mg, 0.6 mmol) and 2-  
methoxyphenylboronic acid (4.62 g, 30.4 mmol). This reaction mixture  
was then heated at 80°C for 12 h, allowed to cool to ambient temperature  
30 then partitioned between ethyl acetate and water. The organics were  
washed with 0.1N sodium hydroxide, water, brine, dried over anhydrous

magnesium sulfate, filtered then pre-adsorbed onto silica. Purification by chromatography on silica gel eluting with dichloromethane:conc. ammonia (99.5:0.5) on a gradient of MeOH (1-4%) afforded 3-(2'-methoxybiphenyl-3-yl)-7-methylimidazo[1,2- $\alpha$ ]pyrimidine as a pale yellow solid (4.4 g, 73%):  $\delta_{\text{H}}$  (400 MHz, DMSO) 2.58 (3H, s), 3.82 (3H, s), 7.03-7.08 (2H, m), 7.16 (1H, d,  $J$  8), 7.38-7.45 (2H, m), 7.53-7.65 (3H, m), 7.74 (1H, s), 7.89 (1H, s), 8.91 (1H, d,  $J$  8);  $m/z$  (ES<sup>+</sup>) 316 (M<sup>+</sup>+H).

### EXAMPLE 17

10

#### 3-(2'-Cyanobiphenyl-3-yl)imidazo[1,2- $\alpha$ ]pyrimidine-7-carboxylic acid methyl ester

4-Ethoxy-2-oxobut-3-enoic acid methyl ester was condensed with 2-aminoimidazole hemisulfate as described in Example 1 to give  
15 imidazo[1,2- $\alpha$ ]pyrimidine-7-carboxylic acid methyl ester as an orange solid:  $\delta_{\text{H}}$  (400 MHz, CDCl<sub>3</sub>) 4.05 (3H, s), 7.68 (1H, d,  $J$  7), 7.72 (1H, d,  $J$  1), 8.06 (1H, d,  $J$  1), 8.59 (1H, d,  $J$  7).

Imidazo[1,2- $\alpha$ ]pyrimidine-7-carboxylic acid methyl ester was brominated as described in Example 1 to give 3-bromoimidazo[1,2- $\alpha$ ]pyrimidine-7-carboxylic acid methyl ester as a cream-coloured solid:  
20  $\delta_{\text{H}}$  (400 MHz, CDCl<sub>3</sub>) 4.07 (3H, s), 7.80 (1H, d,  $J$  7), 8.04 (1H, s), 8.55 (1H, d,  $J$  7).

3-Bromoimidazo[1,2- $\alpha$ ]pyrimidine-7-carboxylic acid methyl ester was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3-(2'-cyanobiphenyl-3-yl)-  
25 imidazo[1,2- $\alpha$ ]pyrimidine-7-carboxylic acid methyl ester as a pale yellow powder:  $\delta_{\text{H}}$  (400 MHz, CDCl<sub>3</sub>) 4.07 (3H, s), 7.50-7.54 (1H, m), 7.58-7.76 (6H, m), 7.81-7.84 (2H, m), 8.18 (1H, s), 9.10 (1H, d,  $J$  7);  $m/z$  (ES<sup>+</sup>) 355 (M<sup>+</sup>+H).

**EXAMPLE 18**3'-(7-Dimethoxymethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile

5           7-Dimethoxymethylimidazo[1,2- $\alpha$ ]pyrimidine was brominated as described in Example 1 to give 3-bromo-7-dimethoxymethylimidazo[1,2- $\alpha$ ]pyrimidine as an off-white solid:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 3.50 (6H, s), 5.28 (1H, s), 7.28 (1H, d,  $J$  7), 7.81 (1H, s), 8.43 (1H, d,  $J$  7);  $m/z$  ( $ES^+$ ) 272/274 ( $M^++H$ ).

10           3-Bromo-7-dimethoxymethylimidazo[1,2- $\alpha$ ]pyrimidine (0.28 g, 1.04 mmol) was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-(7-dimethoxymethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbonitrile (165 mg, 44%) as a yellow powder:  $\delta_H$  (400 MHz,  $CDCl_3$ ) 3.51 (6H, s), 5.30 (1H, s), 7.25 (1H, d,  $J$  7), 7.51 (1H, td,  $J$  8 and 1), 7.57-7.70 (5H, m), 7.76-7.77 (1H, m), 7.82 (1H, dd,  $J$  8 and 1), 7.96 (1H, s), 8.97 (1H, d,  $J$  7);  $m/z$  ( $ES^+$ ) 371 ( $M^++H$ ).

**EXAMPLE 19**3'-[7-([1,2,4]Triazol-1-ylmethyl)imidazo[1,2- $\alpha$ ]pyrimidin-3-yl]biphenyl-2-carbonitrile

20           To a solution of 3'-(7-hydroxymethylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)-biphenyl-2-carbonitrile (0.1 g, 0.31 mmol) in dichloromethane (5 ml) was added carbon tetrabromide (153 mg, 0.46 mmol) and triphenylphosphine (121 mg, 0.46 mmol). This mixture was stirred at ambient temperature for 6 h after which time 1,2,4-triazole sodium salt (84 mg, 0.93 mmol) was added and the reaction stirred at ambient temperature for a further 18 h. The solvent was removed *in vacuo* and the residue purified by silica gel chromatography eluting with dichloromethane on a gradient of methanol (0-10%). Further purification by high performance liquid chromatography

gave 3'-[7-([1,2,4]triazol-1-ylmethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile (15.3 mg):  $\delta_H$  (400 MHz, DMSO) 5.76 (2H, s), 7.32 (1H, d, *J* 7), 7.52-7.84 (8H, m), 8.16 (2H, d, *J* 5), 8.73 (1H, s), 9.22 (1H, s); *m/z* (ES<sup>+</sup>) 378 (M<sup>+</sup>+H).

5

### EXAMPLE 20

#### 3'-(7-Difluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile

4-Ethoxy-1,1-difluorobut-3-en-2-one was condensed with 2-aminoimidazole hemisulfate using the procedure described in Example 1 to give 7-difluoromethylimidazo[1,2-*a*]pyrimidine (16.9 g, 46%) as a pale brown crystalline solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 6.64 (1H, t, *J* 55), 7.26 (1H, d, *J* 7), 7.67 (1H, d, *J* 1), 7.95 (1H, d, *J* 1), 8.60 (1H, d, *J* 7).

7-Difluoromethylimidazo[1,2-*a*]pyrimidine (1.00 g, 5.91 mmol) was brominated as described in Example 1 to give 3-bromo-7-difluoromethylimidazo[1,2-*a*]pyrimidine (0.95 g, 65%) as a white solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 6.67 (1H, t, *J* 55), 7.34 (1H, d, *J* 7), 7.94 (1H, s), 8.57 (1H, d, *J* 7).

3-Bromo-7-difluoromethylimidazo[1,2-*a*]pyrimidine (248 mg, 1.00 mmol) was coupled with 3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 3'-(7-difluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile (201 mg, 58%) as a yellow powder:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 6.67 (1H, t, *J* 55), 7.29 (1H, d, *J* 7), 7.52 (1H, td, *J* 9 and 1), 7.58-7.74 (5H, m), 7.87 (2H, m), 8.01 (1H, s), 9.12 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 347 (M<sup>+</sup>+H).

### EXAMPLE 21

#### 7-Methyl-3-[3-(pyridin-3-yl)phenyl]imidazo[1,2-*a*]pyrimidine

To a degassed solution of trifluoromethanesulfonic acid 3-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)phenyl ester (0.1 g, 0.28 mmol) in 1,4-dioxane

(3 ml) was added potassium phosphate (0.119 g, 0.56 mmol), pyridine-3-boronic acid 1,3-propanediol cyclic ester (0.092 g, 0.56 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.016 g, 13.8  $\mu$ mol) and the mixture heated at 80°C for 22 h. After cooling to ambient temperature the solvent was evaporated *in vacuo*. The residue was stirred with 10% methanol in dichloromethane and the solid material filtered off. The filtrate was applied to two preparative TLC plates (silica gel) and eluted with 6% methanol in dichloromethane. The appropriate band was collected and processed to give an oil which crystallised from diethyl ether as a white solid (26 mg):  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 2.67 (3H, s), 6.80 (1H, d,  $J$  7), 7.42 (1H, td,  $J$  8 and 1), 7.57 (1H, m), 7.65 (2H, m), 7.71 (1H, s), 7.86 (1H, s), 7.92 (1H, m), 8.53 (1H, d,  $J$  7), 7.65 (1H, m), 8.89 (1H, s);  $m/z$  ( $\text{ES}^+$ ) 287 ( $\text{M}^+ + \text{H}$ ).

## EXAMPLE 22

### 7-Methyl-3-[3'-(5-methyl-[1,2,4]oxadiazol-3-yl)biphenyl-3-yl]imidazo[1,2- $\alpha$ ]pyrimidine

To a degassed solution of trifluoromethanesulfonic acid 3-(7-methyl-imidazo[1,2- $\alpha$ ]pyrimidin-3-yl)phenyl ester (2.53 g, 7 mmol) in 1,4-dioxane (25 ml) was added potassium acetate (2.08 g, 21 mmol), bis(pinacolato)diboron (1.97 g, 7.8 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) (0.258 g, 0.35 mmol) and the mixture heated at 80°C for 22 h. After cooling to ambient temperature the reaction was poured into water, extracted with ethyl acetate (3 x 100 ml) and the combined organics were dried over magnesium sulfate. Filtration and evaporation *in vacuo* gave a dark oil which was chromatographed on silica gel, eluting with a gradient of 1 to 10% methanol in dichloromethane. Appropriate fractions were pooled and evaporated to give 7-methyl-3-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]imidazo[1,2- $\alpha$ ]pyrimidine as a cream coloured solid (1.97 g):  $\delta_{\text{H}}$



(400 MHz, CDCl<sub>3</sub>) 1.36 (12H, s), 6.77 (1H, d, *J* 7), 7.52 (1H, m), 7.59 (1H, m), 7.80 (1H, s), 7.87 (1H, m), 7.95 (1H, s), 8.50 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 336 (M<sup>+</sup>+H).

To a degassed solution of 7-methyl-3-[3-(4,4,5,5-tetramethyl-  
5 [1,3,2]dioxaborolan-2-yl)phenyl]imidazo[1,2-*α*]pyrimidine (0.1 g, 0.29 mmol) in 1,4-dioxane (3 ml) was added 3-(3-bromophenyl)-5-methyl-[1,2,4]oxadiazole (0.142 g, 0.59 mmol) (prepared as described in WO 95/27692), potassium phosphate (189 mg, 0.89 mmol) and  
10 tetrakis(triphenylphosphine)palladium(0) (17 mg, 15 μmol) and the mixture heated at 90°C for 18 h. After cooling to ambient temperature the solvent was evaporated *in vacuo* and the residue stirred with 10% methanol in dichloromethane. The solid material was removed by filtration and the filtrate applied to two preparative TLC plates (silica gel) and eluted with 5% methanol in dichloromethane. The appropriate band  
15 was collected and processed, affording a solid which was recrystallised from dichloromethane/ethyl acetate/isohexane to give 7-methyl-3-[3'-(5-methyl-[1,2,4]oxadiazol-3-yl)biphenyl-3-yl]imidazo[1,2-*α*]pyrimidine as a cream coloured solid (32 mg): δ<sub>H</sub> (400 MHz, CDCl<sub>3</sub>) 2.67 (3H, s), 2.68 (3H, s), 6.79 (1H, d, *J* 7), 7.60-7.80 (5H, m), 7.87 (1H, s), 7.92 (1H, m), 8.10 (1H, dd, *J* 7 and 1), 8.32 (1H, s), 8.52 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 368 (M<sup>+</sup>+H).  
20

### EXAMPLE 23

7-Methyl-3-[2'-(3-methyl-[1,2,4]oxadiazol-5-yl)biphenyl-3-yl]imidazo[1,2-  
25 α]pyrimidine

To activated molecular sieves (2.0 g, 4Å powdered) was added ethanol (20 ml) followed by sodium metal (0.435 g, 18.9 mmol), and the mixture heated at reflux for 15 min. After cooling to ambient temperature, acetamide oxime (1.4 g, 18.9 mmol) was added and the  
30 suspension stirred for 15 min. Methyl 2-bromobenzoate (1.32 ml, 9.5 mmol) was added and the mixture heated at reflux for 2 h. The reaction

was poured into water and extracted with ethyl acetate. The organic phase was separated, dried over magnesium sulfate, filtered and evaporated to give a white solid. The crude product was purified by crystallisation from ethyl acetate/isohexane to give 5-(2-bromophenyl)-3-methyl-[1,2,4]oxadiazole as a white solid (0.76 g):  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 2.52 (3H, s), 7.45 (2H, m), 7.76 (1H, dd,  $J$  7.8 and 1.2), 7.97 (1H, dd,  $J$  7.8 and 1.2);  $m/z$  ( $\text{ES}^+$ ) 239:241 (1:1) ( $\text{M}^++\text{H}$ ).

7-Methyl-3-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]-imidazo[1,2- $\alpha$ ]pyrimidine (0.1 g, 0.29 mmol) was coupled to 5-(2-bromophenyl)-3-methyl-[1,2,4]oxadiazole (0.142 g, 0.59 mmol) as described in Example 22 and purified by mass-directed preparative HPLC to give 7-methyl-3-[2'-(3-methyl-[1,2,4]oxadiazol-5-yl)biphenyl-3-yl]imidazo[1,2- $\alpha$ ]pyrimidine as its trifluoroacetate salt (21 mg):  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 2.40 (3H, s), 2.76 (3H, s), 7.07 (1H, d,  $J$  7), 7.43-7.69 (7H, m), 8.03 (1H, s), 8.10 (1H, dd,  $J$  7.8 and 1.1), 8.61 (1H, d,  $J$  7);  $m/z$  ( $\text{ES}^+$ ) 368 ( $\text{M}^++\text{H}$ ).

#### EXAMPLE 24

##### 7-Methyl-3-[3-(thiazol-4-yl)phenyl]imidazo[1,2- $\alpha$ ]pyrimidine

7-Methyl-3-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-imidazo[1,2- $\alpha$ ]pyrimidine (0.1 g, 0.29 mmol) was coupled to 4-bromothiazole (98 mg, 0.59 mmol) as described in Example 22 except using *N,N*-dimethylacetamide in place of 1,4-dioxane as solvent. After cooling to ambient temperature the reaction was poured onto a strong cation exchange cartridge and eluted with methanol. The product was then eluted with a 2.0M solution of ammonia in methanol and evaporated *in vacuo*. The product was purified by mass-directed preparative HPLC to give 7-methyl-3-[3-(thiazol-4-yl)phenyl]imidazo[1,2- $\alpha$ ]pyrimidine as its trifluoroacetate salt (24 mg):  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 2.77 (3H, s), 7.12 (1H, d,  $J$  7), 7.51 (1H, m), 7.64-7.69 (2H, m), 8.05-8.09 (2H, m), 8.17 (1H, m), 8.65 (1H, d,  $J$  7), 8.92 (1H, s);  $m/z$  ( $\text{ES}^+$ ) 293 ( $\text{M}^++\text{H}$ ).

**EXAMPLE 25****3'-(7-Methylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbaldehyde oxime**

5        7-Methyl-3-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]-imidazo[1,2- $\alpha$ ]pyrimidine (0.1 g, 0.29 mmol) was coupled to 2-bromobenzaldehyde oxime (98 mg, 0.59 mmol) as described in Example 24 to give 3'-(7-methylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)biphenyl-2-carbaldehyde oxime as its trifluoroacetate salt (24 mg):  $\delta_H$  (400 MHz,  $d^6$ DMSO) 2.71 (3H, s), 7.42-7.53 (4H, d,  $J$  7), 7.68-7.79 (3H, m), 7.89 (1H, m), 8.01 (1H, s), 8.37  
10 (1H, s), 8.65 (1H, d,  $J$  7), 9.17 (1H, d,  $J$  7.1), 11.36 (1H, bs);  $m/z$  (ES<sup>+</sup>) 329 (M<sup>+</sup>+H).

**EXAMPLE 26****3-[3-(7-Methylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)phenyl]pyridine-2-carbonitrile**

15        7-Methyl-3-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]-imidazo[1,2- $\alpha$ ]pyrimidine (0.1 g, 0.29 mmol) was coupled to 3-bromopyridine-2-carbonitrile (0.11 g, 0.59 mmol) as described in Example  
20 24. The crude product was applied to two preparative plates (silica gel) and eluted with 5% methanol in dichloromethane. The appropriate band was collected and processed, affording a solid which was recrystallised from dichloromethane/ethyl acetate/isohexane to give 3-[3-(7-methylimidazo[1,2- $\alpha$ ]pyrimidin-3-yl)phenyl]pyridine-2-carbonitrile as a  
25 white solid (31 mg):  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 2.67 (3H, s), 6.80 (1H, d,  $J$  7), 7.55 (1H, td,  $J$  8 and 1), 7.63-7.73 (3H, m), 7.77 (1H, m), 7.86 (1H, s), 7.94 (1H, dd,  $J$  8 and 1.6), 8.75 (1H, d,  $J$  1.6), 8.80 (1H, d,  $J$  7);  $m/z$  (ES<sup>+</sup>) 312 (M<sup>+</sup>+H).

**EXAMPLE 27****7-Methyl-3-[3-(pyridin-2-yl)phenyl]imidazo[1,2- $\alpha$ ]pyrimidine**

7-Methyl-3-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]-  
5 imidazo[1,2- $\alpha$ ]pyrimidine (0.1 g, 0.29 mmol) was coupled to 2-  
bromopyridine (55  $\mu$ l, 0.59 mmol) as described in Example 26 to give 7-  
methyl-3-[3-(pyridin-2-yl)phenyl]imidazo[1,2- $\alpha$ ]pyrimidine as a white solid  
(30 mg):  $\delta_H$  (400 MHz,  $CDCl_3$ ) 2.67 (3H, s), 6.79 (1H, d,  $J$  7), 7.30 (1H, m),  
7.55-7.65 (2H, m), 7.77-7.81 (2H, m), 7.87 (1H, s), 8.00 (1H, dd,  $J$  8 and  
10 1.6), 8.19 (1H, d,  $J$  1.5), 8.57 (1H, d,  $J$  7), 8.71 (1H, d,  $J$  1.6);  $m/z$  ( $ES^+$ ) 287  
( $M^+ + H$ ).

**EXAMPLE 28****7-Methyl-3-[3-(thiazol-2-yl)phenyl]imidazo[1,2- $\alpha$ ]pyrimidine**

7-Methyl-3-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]-  
imidazo[1,2- $\alpha$ ]pyrimidine (0.1 g, 0.29 mmol) was coupled to 2-  
bromothiazole (52  $\mu$ l, 0.59 mmol) as described in Example 26 to give 7-  
methyl-3-[3-(pyridin-2-yl)phenyl]imidazo[1,2- $\alpha$ ]pyrimidine as a white solid  
20 (23 mg):  $\delta_H$  (400 MHz,  $CDCl_3$ ) 2.68 (3H, s), 6.81 (1H, d,  $J$  7), 7.39 (1H, d,  $J$   
3.2), 7.57-7.63 (2H, m), 7.87 (1H, s), 7.91 (1H, d,  $J$  3.2), 7.99 (1H, dd,  $J$  8  
and 1.6), 8.17 (1H, m), 8.57 (1H, d,  $J$  7);  $m/z$  ( $ES^+$ ) 293 ( $M^+ + H$ ).

**EXAMPLE 29**

25

**7-Methyl-3-(2'-trifluoromethylbiphenyl-3-yl)imidazo[1,2- $\alpha$ ]pyrimidine**

Trifluoromethanesulfonic acid 3-(7-methylimidazo[1,2- $\alpha$ ]pyrimidin-  
3-yl)phenyl ester (0.1 g, 0.28 mmol) was coupled to 2-  
trifluoromethylbenzenboronic acid (52  $\mu$ l, 0.59 mmol) as described in  
30 Example 21 to give 7-methyl-3-(2'-trifluoromethylbiphenyl-3-yl)-  
imidazo[1,2- $\alpha$ ]pyrimidine as a white solid (35 mg):  $\delta_H$  (400 MHz,  $CDCl_3$ )

2.65 (3H, s), 6.77 (1H, d, *J* 7), 7.39 (1H, m), 7.48-7.62 (6H, m), 7.78 (1H, d, *J* 7.2), 7.82 (1H, s), 8.52 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 354 (M<sup>+</sup>+H).

### EXAMPLE 30

5

#### 3-(2'-Fluorobiphenyl-3-yl)-7-methylimidazo[1,2-*α*]pyrimidine

Trifluoromethanesulfonic acid 3-(7-methylimidazo[1,2-*α*]pyrimidin-3-yl)phenyl ester (0.1 g, 0.28 mmol) was coupled to 2-fluorobenzene-boronic acid (52 μl, 0.59 mmol) as described in Example 21 to give 3-(2'-fluorobiphenyl-3-yl)-7-methylimidazo[1,2-*α*]pyrimidine as a white solid (104 mg): δ<sub>H</sub> (400 MHz, CDCl<sub>3</sub>) 2.67 (3H, s), 6.78 (1H, d, *J* 7), 7.17-7.28 (2H, m), 7.37 (1H, m), 7.46-7.54 (2H, m), 7.60 (2H, m), 7.71 (1H, s), 7.85 (1H, s), 8.56 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 304 (M<sup>+</sup>+H).

15

### EXAMPLE 31

#### 4-Fluoro-3'-(7-methylimidazo[1,2-*α*]pyrimidin-3-yl)biphenyl-2-carbonitrile

7-Methyl-3-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)phenyl]-imidazo[1,2-*α*]pyrimidine (0.2 g, 0.6 mmol) was coupled to 2-bromo-5-fluorobenzonitrile (179 mg, 0.9 mmol) as described in Example 26 to give 4-fluoro-3'-(7-methylimidazo[1,2-*α*]pyrimidin-3-yl)biphenyl-2-carbonitrile as a white solid (85 mg): δ<sub>H</sub> (400 MHz, CDCl<sub>3</sub>) 2.67 (3H, s), 6.83 (1H, d, *J* 7), 7.20 (1H, dt, *J* 8.9 and 3), 7.50-7.70 (6H, m), 7.84 (1H, s), 8.76 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 329 (M<sup>+</sup>+H).

25

### EXAMPLE 32

#### 3-[3-(Imidazol-1-yl)phenyl]-7-trifluoromethylimidazo[1,2-*α*]pyrimidine

To a cooled (-78°C) solution of 3-bromo-7-trifluoromethyl-imidazo[1,2-*α*]pyrimidine (1.0 g, 3.78 mmol) in tetrahydrofuran (20 ml) was added isopropylmagnesium chloride (2.08 ml of a 2.0M solution in

30

tetrahydrofuran, 4.16 mmol). After stirring for 5 min tributyltin chloride (1.2 ml, 4.42 mmol) was added and the reaction stirred for 10 min at -78°C then allowed to warm to ambient temperature to give a solution of 3-tributylstannyl-7-trifluoromethylimidazo[1,2-*a*]pyrimidine in tetrahydrofuran (*ca.* 0.15M): *m/z* (ES<sup>+</sup>) 474, 476, 478 (M<sup>+</sup>+H).

To a degassed solution of 3-tributylstannyl-7-trifluoromethylimidazo[1,2-*a*]pyrimidine (1.4 mmol) was added 1-(3-bromophenyl)-1*H*-imidazole (0.60 g, 2.7 mmol) (prepared by the method of A. Johnson *et al.*, *J. Med. Chem.*, 1969, 12(5), 1024-8) and tetrakis(triphenylphosphine)palladium(0) (218 mg, 0.18 mmol) and the mixture heated at reflux for 18 h. The crude reaction was adsorbed onto silica and chromatographed on silica, eluting on a gradient of 1 to 5% methanol in dichloromethane, to give a yellow oil. Crystallisation from ethyl acetate/isohexane afforded 3-[3-(imidazol-1-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine (0.098 g) as a white solid:  $\delta_{\text{H}}$  (400 MHz; DMSO) 7.15 (1H, s), 7.53 (1H, d, *J* 7.4), 7.73 (1H, dd, *J* 3.1 and 1.6), 7.74 (1H, d, *J* 0.8), 7.78-7.81 (1H, m), 7.89 (1H, t, *J* 1.4), 8.08 (1H, dd, *J* 1.6 and 1.2), 8.39 (2H, d, *J* 7.0), 9.44 (1H, d, *J* 6.7); *m/z* (ES<sup>+</sup>) 330 (M<sup>+</sup>+H).

20

### EXAMPLE 33

#### 3-[3-([1,2,4]Triazol-1-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine

A suspension of 3-bromophenylhydrazine hydrochloride (3.78 g, 17 mmol) in formamide (15 ml) was heated at 140°C for 16 h. The reaction mixture was cooled to ambient temperature, diluted into dichloromethane (100 ml) and washed with water (2 x 100 ml). The organic phase was dried over anhydrous magnesium sulfate, filtered and evaporated to dryness to give 1-(3-bromophenyl)-1*H*-[1,2,4]triazole (3.44 g) as a tan solid:  $\delta_{\text{H}}$  (400 MHz, CDCl<sub>3</sub>) 8.57 (1H, s), 8.11 (1H, s), 7.90 (1H, t, *J* 2), 7.62-7.64 (1H, m), 7.53-7.55 (1H, m), 7.38 (1H, t, *J* 8).

1-(3-Bromophenyl)-1*H*-[1,2,4]triazole (1.7 g, 7.6 mmol) was coupled to 3-tributylstannyl-7-trifluoromethylimidazo[1,2-*α*]pyrimidine (3.8 mmol) by the method of Example 32. Purification by chromatography on silica gel eluting with dichloromethane on a gradient of methanol (1-3%), then  
5 crystallisation from toluene/isohexane, gave 3-[3-([1,2,4]triazol-1-yl)-phenyl]-7-trifluoromethylimidazo[1,2-*α*]pyrimidine as an off-white solid:  $\delta_{\text{H}}$  (400 MHz, DMSO) 9.43 (1H, d, *J* 7), 9.41 (1H, s), 8.38 (1H, s), 8.30 (1H, s), 8.26-8.28 (1H, m), 7.97-8.00 (1H, m), 7.76-7.81 (2H, m), 7.56 (1H, d, *J* 7); *m/z* (ES<sup>+</sup>) 331 (M<sup>++</sup>H).

#### EXAMPLE 34

##### 3-[2'-([1,2,4]Triazol-1-yl)biphenyl-3-yl]-7-trifluoromethylimidazo[1,2-*α*]pyrimidine

15 A suspension of 2-bromophenylhydrazine hydrochloride (5.0 g, 22 mmol) in formamide (10 ml) was heated at 140°C for 16 h. The reaction mixture was cooled to ambient temperature, diluted into dichloromethane (50 ml) and washed with water (3 x 20 ml). The organic phase was dried over anhydrous magnesium sulfate, filtered and evaporated to dryness.

20 Purification by chromatography on silica gel eluting with dichloromethane gave 1-(2-bromophenyl)-1*H*-[1,2,4]triazole (4.80 g) as a tan solid:  $\delta_{\text{H}}$  (400 MHz, CDCl<sub>3</sub>) 8.49 (1H, s), 8.14 (1H, s), 7.76 (1H, dd, *J* 1 and 8), 7.44-7.53 (2H, m), 7.34-7.39 (1H, m).

To a degassed solution of 1-(2-bromophenyl)-1*H*-[1,2,4]triazole (1.0  
25 g, 4.5 mmol) in toluene (50 ml) was added hexabutylditin (4.7 ml, 9.3 mmol), then tetrakis(triphenylphosphine)palladium(0) (160 mg, 139  $\mu$ mol) and the reaction heated at reflux for 48 h. The solvent was evaporated at reduced pressure and the residue chromatographed on silica gel eluting with dichloromethane to give 1-(2-tributylstannylphenyl)-1*H*-

30 [1,2,4]triazole as an oil: *m/z* (ES<sup>+</sup>) 434, 435, 436 (M<sup>++</sup>H).

To a degassed solution of 1-(2-tributylstannylphenyl)-1*H*-[1,2,4]triazole (300 mg, 0.7 mmol) in *N,N*-dimethylformamide (5 ml) was added 1,3-dibromobenzene (0.17 ml, 1.4 mmol), tetrakis(triphenylphosphine)palladium(0) (40 mg, 35  $\mu$ mol), lithium chloride (293 mg, 6.9 mmol), then copper(I) iodide (13 mg, 69  $\mu$ mol), and the mixture was heated at 80°C for 5 h. The reaction was cooled to ambient temperature then poured into water (20 ml) and extracted with dichloromethane (3 x 20 ml). The combined organic phases were dried over anhydrous magnesium sulfate, filtered and evaporated to dryness. Purification by chromatography on silica gel eluting with dichloromethane on a gradient of methanol (0-1%), then by preparative thin-layer chromatography on silica gel using 5% methanol in dichloromethane as eluent, gave 1-(3'-bromobiphenyl-2-yl)-1*H*-[1,2,4]triazole as a colourless gel: *m/z* (ES<sup>+</sup>) 300, 302 (M<sup>+</sup>+H).

1-(3'-Bromobiphenyl-2-yl)-1*H*-[1,2,4]triazole (91 mg, 0.30 mmol) was coupled to 3-tributylstannyl-7-trifluoromethylimidazo[1,2-*a*]pyrimidine (0.25 mmol) by the method of Example 32. The product was obtained by preparative thin-layer chromatography on silica gel with 3% methanol in dichloromethane as eluent followed by high performance liquid chromatography to give 3-[2'-([1,2,4]triazol-1-yl)biphenyl-3-yl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine as a yellow solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 8.57 (1H, d, *J* 7), 7.95-8.11 (3H, m), 7.48-7.66 (7H, m), 7.35 (1H, d, *J* 8), 7.27 (1H, m); *m/z* (ES<sup>+</sup>) 407 (M<sup>+</sup>+H).

### EXAMPLE 35

#### 3'-[7-(Morpholin-4-ylmethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile

To a solution of 3'-(7-hydroxymethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile (0.1 g, 0.31 mmol) in anhydrous dichloromethane under nitrogen was added carbon tetrabromide (153 mg,



0.46 mmol) and triphenylphosphine (121 mg, 0.46 mmol). The resultant mixture was stirred for 6 h after which time morpholine (67  $\mu$ l, 0.77 mmol) was added and the reaction stirred for a further 18 h. The solvent was evaporated *in vacuo* and the residue purified by silica gel chromatography eluting with dichloromethane on a gradient of methanol (0-10%). Further purification by high performance liquid chromatography gave 3'-[7-(morpholin-4-ylmethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile trifluoroacetate salt (18.0 mg):  $\delta_H$  (400 MHz,  $d_6$ DMSO) 3.52 (4H, bs), 4.02 (4H, m), 4.64 (2H, s), 7.50-7.84 (9H, m), 8.12 (1H, bs), 9.17 (1H, s);  $m/z$  (ES<sup>+</sup>) 396 (M<sup>+</sup>+H).

### EXAMPLE 36

#### 4-Fluoro-3'-(7-trifluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile

2-Bromo-5-fluorobenzonitrile and 3-nitrophenylboronic acid were coupled following the procedure in Example 1 to afford 4-fluoro-3'-nitrobiphenyl-2-carbonitrile as a black solid:  $\delta_H$  (360 MHz, CDCl<sub>3</sub>) 7.39-7.48 (2H, m), 7.52-7.64 (1H, m), 7.71 (1H, dd, *J* 8 and 8), 7.89 (1H, d, *J* 8), 8.33-8.37 (2H, m).

4-Fluoro-3'-nitrobiphenyl-2-carbonitrile was reduced by treatment with tin(II) chloride in ethanol and tetrahydrofuran to give 3'-amino-4-fluorobiphenyl-2-carbonitrile as a brown solid:  $\delta_H$  (360 MHz, CDCl<sub>3</sub>) 6.76 (1H, ddd, *J* 8, 2 and 2), 6.80 (1H, dd, *J* 2 and 2), 6.87 (1H, ddd, *J* 8, 1 and 1), 7.27 (1H, dd, *J* 8 and 8), 7.35 (1H, ddd, *J* 8, 8 and 3), 7.41-7.51 (2H, m).

3'-Amino-4-fluorobiphenyl-2-carbonitrile was bromo-deaminated by treatment with 48% hydrobromic acid in 1,4-dioxane, then with sodium nitrite in water at <5°C, then with copper(I) bromide in 48% hydrobromic acid, to give 3'-bromo-4-fluorobiphenyl-2-carbonitrile as a white solid:  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 7.35-7.40 (2H, m), 7.46-7.50 (3H, m), 7.59 (1H, dd, *J* 2 and 1), 7.64 (1H, dd, *J* 2 and 2).

3'-Bromo-4-fluorobiphenyl-2-carbonitrile was converted to 4-fluoro-3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile following the procedure in Example 1 to furnish a brown oil that crystallised on standing:  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 1.36 (12H, s), 7.32-7.37 (1H, m), 7.43-7.54 (3H, m), 7.63-7.68 (1H, m), 7.88-7.90 (2H, m).

3-Bromo-7-trifluoromethylimidazo[1,2-*a*]pyrimidine was coupled with 4-fluoro-3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 4-fluoro-3'-(7-trifluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile (220 mg, 57%) as a yellow powder:  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 7.29 (1H, d, *J* 7), 7.45 (1H, ddd, *J* 8, 1 and 1), 7.53 (1H, dd, *J* 8 and 3), 7.55-7.61 (2H, m), 7.65-7.67 (1H, m), 7.70-7.75 (1H, m), 7.77 (1H, s), 8.14 (1H, s), 9.14 (1H, d, *J* 7);  $m/z$  ( $\text{ES}^+$ ) 383 ( $\text{M}^++\text{H}$ ).

15

### EXAMPLE 37

#### 4-Fluoro-3'-[7-(2-hydroxyprop-2-yl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile

2-(3-Bromoimidazo[1,2-*a*]pyrimidin-7-yl)propan-2-ol was coupled with 4-fluoro-3'-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)biphenyl-2-carbonitrile as described in Example 1 to give 4-fluoro-3'-[7-(2-hydroxyprop-2-yl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile (260 mg, 70%) as a yellow powder:  $\delta_{\text{H}}$  (360 MHz,  $\text{CDCl}_3$ ) 1.61 (6H, s), 4.51 (1H, s), 7.07 (1H, d, *J* 7), 7.45 (1H, ddd, *J* 8, 1 and 1), 7.51-7.60 (3H, m), 7.63-7.69 (2H, m), 7.73 (1H, s), 7.90 (1H, s), 8.94 (1H, d, *J* 7);  $m/z$  ( $\text{ES}^+$ ) 373 ( $\text{M}^++\text{H}$ ).

30

### EXAMPLE 38

#### 3-[3-(Pyridin-3-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine

A mixture of 1,3-dibromobenzene (8.7 g, 36.8 mmol) and pyridine-3-boronic acid-1,3-propanediol cyclic ester (4.0 g, 24.5 mmol) in EtOH (60 ml) and toluene (60 ml) together with 2N Na<sub>2</sub>CO<sub>3</sub> solution (24.5 ml) was degassed with a stream of N<sub>2</sub> for 10 min. Tetrakis(triphenylphosphine)-palladium(0) (0.25 g, 0.22 mmol) was added and the reaction heated at reflux for 14 h. The mixture was concentrated under reduced pressure to remove the organic solvents. The organics were extracted with EtOAc (2 x 125 ml) and then washed with brine (75 ml), dried (MgSO<sub>4</sub>), and concentrated under reduced pressure. The resulting crude residue was purified by column chromatography on silica, using 80% diethyl ether in hexane as the eluent, to yield 3-(3-bromophenyl)pyridine (3.66 g, 64%):  $\delta_{\text{H}}$  (360 MHz, CDCl<sub>3</sub>) 7.30-7.42 (2H, m), 7.47-7.60 (2H, m), 7.72 (1H, s), 7.84 (1H, dt, *J* 8 and 2), 8.62 (1H, dd, *J* 4.8 and 1.5), 8.83 (1H, s).

A mixture of 3-(3-bromophenyl)pyridine (1.65 g, 7.1 mmol), bis-(neopentyl glycolato)diborane (1.75 g, 7.8 mmol), KOAc (2.1 g, 21.2 mmol) and Pd(dppf)Cl<sub>2</sub> (288 mg, 5 mol %) in 1,4-dioxane (60 ml) was degassed with a stream of N<sub>2</sub> for 10 min and then heated at 110°C for 16 h. The reaction mixture was concentrated under reduced pressure and diethyl ether (150 ml) was added. This was extracted with 4N NaOH (3 x 50 ml). These combined basic extracts were neutralised with conc. HCl and then extracted with dichloromethane (3 x 100 ml). The combined organic filtrates were washed with brine (50 ml) and dried (MgSO<sub>4</sub>) to yield 3-[3-(5,5-dimethyl-[1,3,2]dioxaborinan-2-yl)phenyl]pyridine (1.63 mg, 87%):  $\delta_{\text{H}}$  (360 MHz, CDCl<sub>3</sub>)  $\delta$  1.04 (6H, s), 3.80 (4H, s), 7.38 (1H, dd, *J* 7.8 and 4.9), 7.47 (1H, t, *J* 7.8), 7.60-7.68 (1H, m), 7.84 (1H, d, *J* 7.4), 7.96 (1H, dt, *J* 7.8 and 2.0), 8.03 (1H, s), 8.58 (1H, dd, *J* 4.9 and 1.5), 8.85 (1H, s); *m/z* (ES<sup>+</sup>) 267 (M+H<sup>+</sup>).

A mixture of 3-bromo-7-trifluoromethylimidazo[1,2-*a*]pyrimidine (300 mg, 1.12 mmol), the foregoing boronate ester (602 mg, 2.24 mmol), 2N Na<sub>2</sub>CO<sub>3</sub> solution (2.24 ml) and THF (4.5 ml) were degassed with a stream of N<sub>2</sub> for 5 min and then tetrakis(triphenylphosphine)palladium(0) (130

mg, 10 mol %) was added and the reaction was heated at 70°C for 90 min. EtOAc (70 ml) was added and the mixture separated, washed with brine (20 ml), dried (MgSO<sub>4</sub>) and concentrated under reduced pressure while dry loading onto MgSO<sub>4</sub>. The residue was purified by column chromatography on silica using 70% EtOAc in hexanes containing 1% Et<sub>3</sub>N and 1% MeOH as the eluent. The resulting material was taken up in MeOH and poured onto a strong cation exchange cartridge and eluted with methanol. The product was then eluted with 2.0M NH<sub>3</sub> in MeOH and evaporated while dry loading onto silica. Subsequent purification by column chromatography on silica using 2.5% MeOH in dichloromethane containing 1% NH<sub>3</sub> (aq.) gave 3-[3-(pyridin-3-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine (120 mg, 31%):  $\delta_H$  (360 MHz, d<sub>6</sub>-DMSO) 7.48-7.58 (2H, m), 7.73 (1H, t, *J* 7.7), 7.81 (1H, d, *J* 7.7), 7.87 (1H, d, *J* 7.7), 8.12 (1H, s), 8.21 (1H, dt, *J* 8.1 and 2.0), 8.37 (1H, s), 8.62 (1H, dd, *J* 4.8 and 1.5), 9.03 (1H, d, *J* 2), 9.42 (1H, d, *J* 7.2); *m/z* (ES<sup>+</sup>) 341 (M<sup>+</sup>).

### EXAMPLE 39

#### 3-[3-([1,2,4]Triazol-4-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine

A mixture of 1,2-bis[(dimethylamino)methylene]hydrazine dihydrochloride (6.25 g, 29.0 mmol) and 3-bromoaniline (5.0 g, 29.0 mmol) in toluene (100 ml) was heated at reflux for 14 h. The resulting mixture was cooled to room temperature and H<sub>2</sub>O (100 ml) added. The layers were separated and the organics were washed with brine (75 ml), dried (MgSO<sub>4</sub>), and concentrated under reduced pressure. The crude residue was purified by column chromatography on silica, using 2% MeOH in EtOAc as the eluent, to yield 4-(3-bromophenyl)-4*H*-[1,2,4]triazole (3.33 g, 51%):  $\delta_H$  (360 MHz, CDCl<sub>3</sub>) 7.32-7.38 (1H, m), 7.43 (1H, t, *J* 8), 7.59 (1H, s), 7.63 (2H, d, *J* 8), 8.47 (2H, s); *m/z* (ES<sup>+</sup>) 223, 225 (1:1) (M<sup>+</sup>).


4-(3-Bromophenyl)-4*H*-[1,2,4]triazole (2.5 g, 11.1 mmol), bis(neopentyl glycolato)diborane (2.77 g, 12.2 mmol), KOAc (3.3 g, 33.5

mmol) and Pd(dppf)Cl<sub>2</sub> (456 mg, 5 mol %) in 1,4-dioxane (100 ml) were reacted as described in Example 38 to yield 4-[3-(5,5-dimethyl-[1,3,2]dioxaborinan-2-yl)phenyl]-4*H*-[1,2,4]triazole (0.78 g, 27%):  $\delta_H$  (360 MHz, CDCl<sub>3</sub>) 1.04 (6H, s), 3.80 (4H, s), 7.40-7.50 (1H, m), 7.53 (1H, *J* 7.5), 7.80 (1H, s), 7.89 (1H, d, *J* 7.5), 8.57 (2H, s); *m/z* (ES<sup>+</sup>) 257 (M<sup>+</sup>+H).

A mixture of 3-bromo-7-trifluoromethylimidazo[1,2-*a*]pyrimidine (139 mg, 0.53 mmol), the foregoing boronate ester (269 mg, 1.05 mmol) and K<sub>3</sub>PO<sub>4</sub> (483 mg, 2.1 mmol) in DMA (3 ml) was degassed with a stream of N<sub>2</sub> for 5 min and then tetrakis(triphenylphosphine)palladium(0) (60 mg, 10 mol %) was added and the reaction heated at 65°C for 70 min. EtOAc (100 ml) was added and the mixture washed with H<sub>2</sub>O (3 x 100 ml) and brine (100 ml), dried (MgSO<sub>4</sub>) and concentrated under reduced pressure while dry loading onto silica. The residue was purified by column chromatography on silica using 5% MeOH in dichloromethane to yield 3-[3-([1,2,4]triazol-4-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine (70 mg, 40%), which was then recrystallised from EtOAc/isohexanes:  $\delta_H$  (400 MHz, d<sub>6</sub>-DMSO) 7.55 (1H, d, *J* 7.2), 7.70-7.90 (3H, m), 8.16 (1H, s), 8.39 (1H, s), 9.24 (2H, s), 9.48 (1H, d, *J* 7.2); *m/z* (ES<sup>+</sup>) 331 (M<sup>+</sup>+H).

**CLAIMS:**

1. A compound of formula I, or a salt or prodrug thereof:



(I)

wherein


Y represents a chemical bond, an oxygen atom, or a -NH- linkage;

Z represents an optionally substituted aryl or heteroaryl group;

10 R<sup>1</sup> represents hydrogen, hydrocarbon, a heterocyclic group, halogen, cyano, trifluoromethyl, nitro, -OR<sup>a</sup>, -SR<sup>a</sup>, -SOR<sup>a</sup>, -SO<sub>2</sub>R<sup>a</sup>, -SO<sub>2</sub>NR<sup>a</sup>R<sup>b</sup>, -NR<sup>a</sup>R<sup>b</sup>, -NR<sup>a</sup>COR<sup>b</sup>, -NR<sup>a</sup>CO<sub>2</sub>R<sup>b</sup>, -COR<sup>a</sup>, -CO<sub>2</sub>R<sup>a</sup>, -CONR<sup>a</sup>R<sup>b</sup> or -CR<sup>a</sup>=NOR<sup>b</sup>; and

15 R<sup>a</sup> and R<sup>b</sup> independently represent hydrogen, hydrocarbon or a heterocyclic group.

2. A compound as claimed in claim 1 represented by formula IIA, and salts and prodrugs thereof:



(IIA)

wherein


Z is as defined in claim 1;

5         $R^{11}$  represents hydrogen,  $C_{1-6}$  alkyl, halo( $C_{1-6}$ )alkyl, dihalo( $C_{1-6}$ )alkyl, hydroxy( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkoxy( $C_{1-6}$ )alkyl, di( $C_{1-6}$ )alkoxy( $C_{1-6}$ )alkyl,  $C_{3-7}$  cycloalkyl,  $C_{3-7}$  heterocycloalkyl( $C_{1-6}$ )alkyl, heteroaryl,  $C_{1-6}$  alkyl-heteroaryl, heteroaryl( $C_{1-6}$ )alkyl, halogen, cyano, trifluoromethyl,  $C_{1-6}$  alkoxy, formyl,  $C_{2-6}$  alkylcarbonyl,  $C_{2-6}$  alkoxy carbonyl or  $-CR^4=NOR^5$ ;

10         $R^4$  represents hydrogen or  $C_{1-6}$  alkyl; and

$R^5$  represents hydrogen,  $C_{1-6}$  alkyl, hydroxy( $C_{1-6}$ )alkyl or di( $C_{1-6}$ )alkylamino( $C_{1-6}$ )alkyl.


3.        A compound as claimed in claim 2 represented by formula  
15        IIB, and salts and prodrugs thereof:



(IIB)

wherein Z is as defined in claim 1.

4. A compound as claimed in claim 2 represented by formula IIC, and salts and prodrugs thereof:



5

(IIC)

wherein

X represents CH or N;

10 R<sup>6</sup> represents fluoro, cyano, trifluoromethyl, methoxy, methyloxadiazolyl, triazolyl or -CR<sup>2</sup>=NOR<sup>3</sup>;

R<sup>7</sup> represents hydrogen or fluoro;

R<sup>2</sup> represents hydrogen or methyl;


R<sup>3</sup> represents hydrogen, hydroxyethyl or dimethylaminoethyl; and

R<sup>11</sup> is as defined in claim 2.

15

5. A compound as claimed in claim 2 represented by formula IID, and salts and prodrugs thereof:





(IID)

wherein

W represents CH or N; and

5 R<sup>11</sup> is as defined in claim 2.

6. A compound selected from:

3'-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

and salts and prodrugs thereof.

10

7. A compound selected from:

3'-(imidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-trifluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-[7-(1,1-dimethoxyethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-

15 carbonitrile;

3'-(7-acetylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-isopropylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-cyclopropylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-*tert*-butylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

20 3'-(7-cyclobutylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-methoxyimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-hydroxymethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-fluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

3'-(7-formylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

- 3'-(7-hydroxyiminomethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;
- 3-(2'-cyanobiphenyl-3-yl)imidazo[1,2-*a*]pyrimidine-7-carbonitrile;
- 3-(2'-methoxybiphenyl-3-yl)-7-methylimidazo[1,2-*a*]pyrimidine;
- 5 3-(2'-cyanobiphenyl-3-yl)imidazo[1,2-*a*]pyrimidine-7-carboxylic acid methyl ester;
- 3'-(7-dimethoxymethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;
- 3'-[7-([1,2,4]triazol-1-ylmethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile;
- 10 3'-(7-difluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;
- 7-methyl-3-[3-(pyridin-3-yl)phenyl]imidazo[1,2-*a*]pyrimidine;
- 7-methyl-3-[3'-(5-methyl-[1,2,4]oxadiazol-3-yl)biphenyl-3-yl]imidazo[1,2-*a*]pyrimidine;
- 15 7-methyl-3-[2'-(3-methyl-[1,2,4]oxadiazol-5-yl)biphenyl-3-yl]imidazo[1,2-*a*]pyrimidine;
- 7-methyl-3-[3-(thiazol-4-yl)phenyl]imidazo[1,2-*a*]pyrimidine;
- 3'-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbaldehyde oxime;
- 3-[3-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)phenyl]pyridine-2-carbonitrile;
- 20 7-methyl-3-[3-(pyridin-2-yl)phenyl]imidazo[1,2-*a*]pyrimidine;
- 7-methyl-3-[3-(thiazol-2-yl)phenyl]imidazo[1,2-*a*]pyrimidine;
- 7-methyl-3-(2'-trifluoromethylbiphenyl-3-yl)imidazo[1,2-*a*]pyrimidine;
- 3-(2'-fluorobiphenyl-3-yl)-7-methylimidazo[1,2-*a*]pyrimidine;
- 4-fluoro-3'-(7-methylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;
- 25 3-[3-(imidazol-1-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine;
- 3-[3-([1,2,4]triazol-1-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine;
- 3-[2'-([1,2,4]triazol-1-yl)biphenyl-3-yl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine;
- 3'-[7-(morpholin-4-ylmethyl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile;
- 30 and salts and prodrugs thereof.

8. A compound selected from:

4-fluoro-3'-(7-trifluoromethylimidazo[1,2-*a*]pyrimidin-3-yl)biphenyl-2-carbonitrile;

5 4-fluoro-3'-[7-(2-hydroxyprop-2-yl)imidazo[1,2-*a*]pyrimidin-3-yl]biphenyl-2-carbonitrile;

3-[3-(pyridin-3-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine;

3-[3-([1,2,4]triazol-4-yl)phenyl]-7-trifluoromethylimidazo[1,2-*a*]pyrimidine;

and salts and prodrugs thereof.

10

9. A pharmaceutical composition comprising a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof or a prodrug thereof in association with a pharmaceutically acceptable carrier.

15


10. The use of a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof or a prodrug thereof for the manufacture of a medicament for the treatment and/or prevention of adverse neurological conditions.

20


11. A process for the preparation of a compound as claimed in claim 1, which comprises:

(A) reacting a compound of formula III with a compound of formula IV:

25




(III)




(IV)

wherein Y, Z and R<sup>1</sup> are as defined in claim 1, L<sup>1</sup> represents a suitable leaving group, and M<sup>1</sup> represents a boronic acid moiety -B(OH)<sub>2</sub> or a cyclic ester thereof formed with an organic diol, or M<sup>1</sup> represents -Sn(Alk)<sub>3</sub> in which Alk represents a C<sub>1-6</sub> alkyl group; in the presence of a transition metal catalyst; or

(B) reacting a compound of formula V with a compound of formula VI:




(V)




(VI)

wherein Y, Z and R<sup>1</sup> are as defined in claim 1, and L<sup>1</sup> and M<sup>1</sup> are as defined above; in the presence of a transition metal catalyst; or

(C) reacting a compound of formula VII with a compound of formula VIII:



(VII)




(VIII)


wherein Z and R<sup>1</sup> are as defined in claim 1, and L<sup>1</sup> and M<sup>1</sup> are as defined above; in the presence of a transition metal catalyst; or

(D) reacting a compound of formula IX with a compound of formula

X:




(IX)



(X)

wherein Z and R¹ are as defined in claim 1, and L¹ and M¹ are as defined above; in the presence of a transition metal catalyst; or


- 5 (E) reacting a compound of formula X as defined above with a compound of formula XI:



(XI)

- 10 wherein R¹ is as defined in claim 1; or

(F) reacting a compound of formula X as defined above with a compound of formula XII:




(XII)


wherein  $R^1$  is as defined in claim 1; or

(G) reacting a compound of formula XVI with a compound of formula XVII:

5




(XVI)




(XVII)

wherein Y, Z and  $R^1$  are as defined in claim 1, and  $L^3$  represents a suitable leaving group; or

10 (H) reacting a compound of formula XXI with a compound of formula XXII:



(XXI)



(XXII)

15 wherein Y and Z are as defined in claim 1,  $R^{1a}$  represents an aryl or heteroaryl moiety, and  $L^4$  represents a suitable leaving group; in the presence of a transition metal catalyst; and

(J) if desired, converting a compound of formula I initially obtained into a further compound of formula I by standard methods.

20

12. A method for the treatment and/or prevention of adverse neurological conditions which comprises administering to a patient in need of such treatment an effective amount of a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof or a  
5 prodrug thereof.

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/GB 01/02158A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07D487/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 19323 A (TEUBER LENE ;WAETJEN FRANK (DK); NEUROSEARCH AS (DK); ICHIMARU YAS) 22 April 1999 (1999-04-22) the whole document	1-12
Y	WO 96 33191 A (NEUROSEARCH AS ;MEIJI SEIKA KAISHA (JP); TEUBER LENE (DK); WAETJEN) 24 October 1996 (1996-10-24) the whole document	1-12
Y	EP 0 616 807 A (NEUROSEARCH AS ;MEIJI SEIKA KAISHA (JP)) 28 September 1994 (1994-09-28) the whole document	1-12
Y	EP 0 563 001 A (NEUROSEARCH AS) 29 September 1993 (1993-09-29) the whole document	1-12
	--- -/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

10 August 2001

Date of mailing of the international search report

23/08/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Stellmach, J



## INTERNATIONAL SEARCH REPORT

In ternational Application No  
PCT/GB 01/02158

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 36423 A (MERCK SHARP & DOHME ;MOORE KEVIN WILLIAM (GB); CARLING WILLIAM ROB) 22 July 1999 (1999-07-22) the whole document	1-12
Y	--- MARTIN I L ET AL: "BENZODIAZEPINE RECOGNITION SITE LIGANDS AND GABAA RECEPTORS" EXPERT OPINION ON THERAPEUTIC PATENTS, ASHLEY PUBLICATIONS, GB, vol. 9, no. 10, 1999, pages 1347-1358, XP001007805 ISSN: 1354-3776 the whole document	1-12
Y	--- TEUBER L ET AL: "LIGANDS FOR THE BENZODIAZEPINE BINDING SITE - A SURVEY" CURRENT PHARMACEUTICAL DESIGN, BENTHAM SCIENCE PUBLISHERS, SCHIPHOL, NL, vol. 5, 1999, pages 317-343, XP001008419 ISSN: 1381-6128 the whole document	1-12
Y	--- TULLY, R. ET AL.: "General Approach Leading to the Development of Imidazoquinoline and Imidazopyrimidine Benzodiazepine Receptor Ligands" DRUG DEVELOPMENT RES., vol. 22, 1991, pages 299-308, XP001013094 NEW YORK the whole document	1-12
Y	--- ZHANG P ET AL: "STUDIES OF NOVEL IMIDAZOBENZODIAZEPINE LIGANDS AT GABAA/BZR SUBTYPES: EFFECT OF C(3) SUBSTITUENTS ON RECEPTOR SUBSITE SELECTIVITY" MEDICINAL CHEMISTRY RESEARCH, BIRKHAUSER, BOSTON, US, vol. 5, no. 7, 1995, pages 487-495, XP000569544 ISSN: 1054-2523 the whole document	1-12
Y	--- SIEGHART W: "STRUCTURE AND PHARMACOLOGY OF GAMMA-AMINOBUTYRIC ACID RECEPTOR SUBTYPES" PHARMACOLOGICAL REVIEWS, WILLIAMS AND WILKINS INC., BALTIMORE, MD., US, vol. 47, no. 2, 1995, pages 181-234, XP001008358 ISSN: 0031-6997 the whole document	1-12

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/02158

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9919323 A	22-04-1999	AU 9433298 A CN 1275127 T EP 1023286 A TW 424088 B ZA 9809238 A	03-05-1999 29-11-2000 02-08-2000 01-03-2001 17-05-1999
WO 9633191 A	24-10-1996	AT 194836 T AU 5501496 A AU 695957 B AU 5689196 A AU 699623 B AU 5690696 A BR 9608048 A BR 9608056 A CA 2217601 A CA 2218493 A CA 2218552 A CN 1182427 A CN 1182426 A CN 1182425 A CZ 9703291 A CZ 9703292 A DE 69609413 D DE 69609413 T DK 821682 T EE 9700282 A EE 9700283 A WO 9633194 A WO 9633192 A EP 0821683 A EP 0821684 A EP 0821682 A ES 2150671 T GR 3034558 T HU 9801692 A HU 9802272 A JP 11501320 T JP 11511734 T JP 3040485 B JP 11501321 T NO 974843 A NO 974844 A NZ 307521 A NZ 307532 A PL 322892 A PL 322944 A PT 821682 T RU 2135493 C RU 2136676 C SK 139997 A SK 140697 A TR 9701218 T TR 9701219 T US 5902813 A US 5922724 A US 5922725 A	15-08-2000 07-11-1996 27-08-1998 07-11-1996 10-12-1998 07-11-1996 30-11-1999 30-11-1999 24-10-1996 24-10-1996 24-10-1996 20-05-1998 20-05-1998 20-05-1998 18-03-1998 18-03-1998 24-08-2000 23-11-2000 04-12-2000 15-06-1998 15-06-1998 24-10-1996 24-10-1996 04-02-1998 04-02-1998 04-02-1998 01-12-2000 31-01-2001 29-03-1999 28-09-1999 02-02-1999 12-10-1999 15-05-2000 02-02-1999 15-12-1997 16-12-1997 29-04-1999 29-03-1999 02-03-1998 02-03-1998 30-11-2000 27-08-1999 10-09-1999 06-05-1998 06-05-1998 21-08-1998 21-03-1998 11-05-1999 13-07-1999 13-07-1999
EP 0616807 A	28-09-1994	AT 168007 T	15-07-1998

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/02158

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0616807 A		AU 675484 B	06-02-1997
		AU 5752194 A	29-09-1994
		CA 2119511 A	25-09-1994
		CN 1099391 A,B	01-03-1995
		DE 69411424 D	13-08-1998
		DE 69411424 T	28-01-1999
		DK 616807 T	19-04-1999
		ES 2119124 T	01-10-1998
		FI 941378 A	25-09-1994
		JP 7002838 A	06-01-1995
		NO 941052 A	26-09-1994
		NZ 260050 A	26-01-1996
		US 5554632 A	10-09-1996
		US 5554630 A	10-09-1996
		ZA 9402079 A	24-10-1994
EP 0563001 A	29-09-1993	AT 134621 T	15-03-1996
		AU 3540693 A	30-09-1993
		CA 2092211 A	27-09-1993
		DE 69301618 D	04-04-1996
		DE 69301618 T	22-08-1996
		DK 563001 T	05-08-1996
		ES 2085133 T	16-05-1996
		FI 931305 A	27-09-1993
		GR 3019204 T	30-06-1996
		JP 6049037 A	22-02-1994
		NO 303779 B	31-08-1998
		NZ 247233 A	28-03-1995
		US 5360809 A	01-11-1994
		ZA 9302090 A	15-10-1993
WO 9936423 A	22-07-1999	AU 1977799 A	02-08-1999
		EP 1047697 A	02-11-2000

